arXiv:1305.0255v1 [math.DG] 1 May 2013

Bessel Functions, Heat Kernel and the Conical Kähler-Ricci Flow.

Xiuxiong Chen, Yuanqi Wang

May 2, 2013

Abstract

Following Donaldson's oppenness theorem on deforming a conical Kähler-Einstein metric, we prove a parabolic Schauder-type estimate with respect to conical metrics. As a corollary, we show that the conical Kähler-Ricci Flow exists for short time. The key is to establish the relevant heat kernel estimates, where we use the Weber's formula on Bessel function of the second kind and Carslaw's heat kernel representation in [7].

1 Introduction.

Let $(M,[\acute{\omega}])$ be a compact Kähler manifold with a smooth Kähler metric $\acute{\omega}$. Let D be a smooth divisor in M. Let L be the line bundle associated to D and let S be the holomorphic section which defines D. Following Donaldson[17], we want to study Kähler metrics on $M\setminus D$ with cone singularities of cone angle $2\beta\pi$ transverse to D, where $0<\beta<1$. In [17], Donaldson studied the linear theory of how conical Kähler-Einstein metrics deform when the cone angle varies. In particular, he proved that the set of cone angles where M admits a Kähler-Einstein metric is open if the only holomorphic vector field tangent to D is the zero vector field. This "openness theorem" fits well into an ambitious program [18] of Donaldson in which he proposed a new continuity approach to attack the renown Kähler-Einstein problem via deforming cone angles. Namely, first show the existence of Kähler-Einstein metrics when conical angle is arbitrary small. Then, gradually open up the angle and show this process is both open and closed, using stability conditions. Donaldson's work [17], as well as his program, inspired a lot new research activities surrounding the existence of conical Kähler -Einstein metrics. The later topic (Singular Kähler Einstein metrics) really goes back much earlier in history (c.f., Yau [51], Tian-Yau, [41][42], etc). For more recent references, we refer to two recent works [5] and [19] and reference therein.

In this paper, following Donaldson's work on the green function in [17], we study the existence of an evolution process. On one hand, we want to deform this Kähler metric in the negative direction of its Ricci form; on the other hand, we want to keep the cone structure fixed during the"evolution" process. The second goal poses the main analytic challenge. A crucial estimate in [17] is the estimate of the Green function for the standard flat conical metric. It is somewhat surprising that the proof of such an estimate relies heavily on direct calculations and estimate of special functions (Bessel functions). In our approach of evolution process, inevitably we need to estimate the heat Kernel for conical metrics. As expected, such an estimate need to employ heavy calculation on Bessel functions. The main conclusion of ours are summarized in Theorem 1.2, 1.8, 1.11. The diagram in Figure 1 is an overall description of the organization of this article.

It is perhaps worthwhile to give a brief account to the history of the Kähler- Einstein metric problem first. In 1950s, E. Calabi asked a famous question: if the first Chern class $C_1 < 0, = 0, > 0$, does there exist a Kähler-Einstein metric with Ricci curvature $< 0, = 0, > 0$ respectively? The case of $C_1 < 0$ is settled by Aubin and by Yau , while Yau solved the case $C_1 = 0$. For the case $C_1 > 0$, among work of others, Tian give a complete solution to the existence of KE metric on Fano surfaces. In higher dimensions, the existence problem is very hard. In early 1980s, Yau conjectured that the existence of Kähler-Einstein metric is related to certain algebraic notions of "stability". Tian [40] introduced a notion called K-stability and proved that it is a necessary condition for the existence of KE metrics. The K-stability was reformulated later in more algebraic ways by Donaldson [16]. The necessary part has been established by Tian[40], Stoppa[35], Berman[2] with various generalities. The existence part is much more difficult. Recently, Chen-Donaldson-Sun confirmed this conjecture of Yau in a series of work [9], [10],[11], [12].

While the main goal stated in Donaldson's program has been successfully tackled, it opens a "door" for exciting future study in Kähler geometry. Following Calabi, one might ask if $(M, (1-\beta)D)$ supports a Kähler-Einstein metric with cone angle $2\beta\pi$ transversal to D. This existence problem can certainly be reduced to a complex Monge-Ampere equation in M and one may attempt to solve its existence via the continuous method. This approach has been taken by a number of authors. As examples, we list a few: the work of Berman [1], Brendle[3] Jeffres-Mazzeo-Rubinstein [24], Li-Sun [26], Song-Wang[34], and Campana- Guenancia-Paun [5]. Calamai -Zheng studied the geodesics in the space of Kähler cone metrics in [4] and A. Hugh [21] studied the geodesic in the space of Kähler cusp metrics. Historically, existence and uniqueness of conical KE metric over Riemann surfaces has been extensively studied. See Troyanov's work [46] and McOwen's work [29] for the existence, and Luo-Tian's work [28] for the uniqueness.

First, let us define a conical Kähler metric.

Definition 1.1. *For any $\alpha \in (0, \min\{\frac{1}{\beta}-1, 1\})$, a Kähler form ω is said to be an (α, β) conical Kähler metric on $(M, (1-\beta)D)$ if it satisfies the following conditions*

1. *ω is a closed positive $(1,1)$ current on M.*
2. *For any point $p \in D$, there exists a small holomorphic chart $(\mathcal{U}, \{z_i\})$ such that in this chart, ω is quasi conformal to the standard cone metric*
$$\frac{\sqrt{-1}}{2}|z_1|^{2\beta-2} dz_1 \wedge d\bar{z}_1 + \frac{\sqrt{-1}}{2}\sum_{j=2}^{n} dz_j \wedge d\bar{z}_j.$$
3. *There exists a $\phi \in C^{2,\alpha,\beta}(M)$ and a smooth Kähler metric $\acute{\omega}$ such that*
$$\omega = \acute{\omega} + i\partial\bar{\partial}\phi.$$
See Definition 2.2 for the definition of the function space $C^{2,\alpha,\beta}(M)$.

Notice that the following model metric defined in [17] satisfies the above definition.

$$\omega_D = \omega^{'} + \delta i\partial\bar{\partial}|S|^{2\beta}, \tag{1}$$

$\omega^{'}$ is a smooth Kähler metric over M and δ is sufficiently small. Similar to (4), ω_D is a (α_0, β) metric, $\alpha_0 = \min\{\frac{2}{\beta} - 2,\ 1\}$.

Now we are ready to introduce the conical Kähler-Ricci flow precisely as

$$\frac{\partial \omega_g}{\partial t} = \beta\omega_g - Ric(g) + 2\pi(1-\beta)[D], \tag{2}$$

where the initial metric $g(0)$ is an (α, β) type Kähler metric in M. (2) should be considered as an equaiton of closed currents over the whole M (not only over $M\backslash D$). Over $M\backslash D$, (2) reduces exactly to the usual Ricci flow equation:

$$\frac{\partial \omega_g}{\partial t} = \beta\omega_g - Ric(g).$$

We can also consider more general conical Kähler-Ricci flows as

$$\frac{\partial \omega_g}{\partial t} = \mu\omega_g - Ric(g) + 2\pi(1-\beta)[D].$$

for any number μ, and the short time existence as Theorem 1.2 holds equally well. For the sake of brevity and to make the reader understand better, we only consider the flow (2) in this article.

At the level of potentials, we have

$$\frac{\partial \phi}{\partial t} = \log \frac{(\omega_D + \sqrt{-1}\partial\bar{\partial}\phi)^n}{\omega_D^n} + \beta\phi + h_{\omega_D}. \tag{3}$$

Here h_{ω_D} satisfies $\sqrt{-1}\partial\bar{\partial}h_{\omega_D} = \beta\omega_D - Ric(\omega_D) + 2\pi(1-\beta)[D]$ over M. The h_{ω_D} has a nice expression as

$$h_{\omega_D} = \beta\delta|S|^{2\beta} + \log\frac{\omega_D^n|S|^{2-2\beta}}{\omega^n} + \acute{F},$$

where $\acute{F}$ is a smooth function over M. Routine calculation shows that

$$h_{\omega_D} \in C^{\alpha_0,\beta},\ \alpha_0 = \min\{\frac{2}{\beta} - 2,\ 1\}. \tag{4}$$

The main problem in this paper is to show the following: if the initial metric ω_0 is of (α,β) type, do we have a one parameter family of (α,β) type conical Kähler metrics $\{\omega(t)(t \in [0,T])\}$ initiated from ω_0 which satisfies the Ricci flow equation? Unlike the classical settings, the short time existence of such a flow is a real challenge: the nature of heat flows is to "smooth" singularities, while the key point in this flow is to preserve the singularity structure.

Our main theorem on the short time existence of conical Kähler-Ricci flow can be formulated as follows:

Theorem 1.2. *Suppose g_0 is an $(\acute{\alpha},\beta)$ conical Kähler metric in $(M,(1-\beta)D)$ where $\alpha' \in (0,\min\{\frac{1}{\beta}-1,1\})$. For any $\alpha \in (0,\alpha')$, there exists an $T_0 > 0$ (which depends on g_0) such that the conical Kähler-Ricci flow (2) initiated from g_0 admits a solution $g(t)$, $t \in [0,T_0]$ which is smooth (both in space and time) in $(M \setminus D) \times [0,T_0]$. Moreover we have*

- *for every $t \in [0,T_0]$, $g(t)$ is an (α,β) conical metric in $(M,(1-\beta)D)$;*
- *$g(t)$ is a $C^{\alpha,\frac{\alpha}{2},\beta}[0,T_0]$-family of conical metrics (see Definition 2.5).*

- *for any* $0 < \widehat{\alpha} \leq \alpha$, $g(t)$ *is the unique solution of (2) in the class of* $C^{\widehat{\alpha}, \frac{\widehat{\alpha}}{2}, \beta}[0, T]$*-family of metrics.*

Remark 1.3. For short time existence theorem, we allow D to be a reducible, smooth divisor. In [17], one needs the kernel of the linearized operator to be trivial to show that the angle can be perturbed. This is not needed in our case since we don't change angles. We only need the Schauder estimates for Linearized operator (c.f. Theorem 1.7 below).

In fact, a more general version is true (see Theorem 2.7). In Theorem 2.7, we allow the angles on each component of D to be different.

Remark 1.4. When $n = 1$ the short time existence in a different function space is proved by Yin in 2009 in [52], using a very different method from ours. Recently, Mazzeo announced another short time existence theorem on conical Ricci flows over conical Riemann surfaces, which is proved in an unpulished work jointly by Mazzeo, Rubinstein, and Sesum. Both results are very interesting to us and are relevant to the dimension 2 (complex dimension 1) part of our work.

Remark 1.5. It's interesting to fit Theorem 1.2 into a more general picture. To be precise, we have at least three kinds of Ricci flows starting from an (α, β) metric as follows. The first kind is as the solution in Theorem 1.2 (c.f. the preceding Remark), which perserves the cone structure; the second kind is as the solutions studied by Topping over incomplete Riemann surfaces (cf. [47]), which flow incomplete (2-dimensional) metrics instaneously to be complete; the third kind is as the solutions considered by Chen-Tian-Zhang in [14] (and Chen-Ding in [8]), which smooth out the conical singularities immediately after the flows start.

Remark 1.6. When $\beta = 1$, this reduces to the renown Kähler-Ricci flow. After the fundamental work of G. Perelman in the Kähler-Ricci flow, this has become a powerful tool in attacking the existence of Kähler-Einstein problems in Fano manifolds. There has been extensive research done in this subject. We refer interested readers to [45], [13], [36], and references therein for further readings.

In a sequel of this paper [15], we will prove the long time existence of this conical Kähler-Ricci flow. As consequence, we prove the convergence of the CKRF when the modified first Chern class is either negative or vanishes (cf. Cao[6] for comparison in the smooth KRF settings). We will also study the extension of Perelman's functional to this flow.

At the level of potential functions, Theorem 1.2 is implied by the following theorem on parabolic Monge-Ampere equations which we use all efforts to prove.

Theorem 1.7. *Assumptions as in Theorem 1.2. Suppose* $0 < T < \infty$ *and* $f \in C^{2+\acute{\alpha},1+\frac{\acute{\alpha}}{2},\beta}[0,T]$. *Suppose* μ *is a constant. Then there exists an* $0 < T_0 \leq T$ *such that the conical Kähler-Ricci flow equation*

$$\begin{cases} \frac{\partial \phi}{\partial t} = \log \frac{(\omega_0+\sqrt{-1}\partial\bar{\partial}\phi)^n}{\omega_0^n} + \mu\phi + f \\ \phi = 0 \ \textit{when} \ t = 0 \end{cases}$$

admits a solution $\phi \in C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T_0]$. *Furthermore, for any* $0 < \widehat{\alpha} \leq \alpha$, ϕ *is the unique solution in* $C^{2+\widehat{\alpha},1+\frac{\widehat{\alpha}}{2},\beta}[0,T_0]$.

The crucial ingredients to establish Theorem 1.7 are the following Schauder type estimates.

Theorem 1.8. *Suppose* $0 < T < \infty$ *and* $\{a(t),\ t \in [0,T]\}$ *is a* $C^{\alpha,\frac{\alpha}{2},\beta}[0,T]$ *family of* (α,β) *metrics (See Definition 2.5). Then there exists a constant* C_a *(depending on* $a(t)$*) with the following property. For any* $v \in C^{\alpha,\frac{\alpha}{2},\beta}[0,T]$, *the parabolic equation*

$$\frac{\partial u}{\partial t} = \Delta_{a(t)}u + v,\ \ u(0,x) = 0 \tag{5}$$

admits a unique solution u *in* $C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$. *Moreover, the following estimates hold.*

$$|u|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \leq C_a|v|_{\alpha,\frac{\alpha}{2},\beta,M\times[0,T]};$$
$$|u|_{2,\alpha,\beta,M\times[0,T]} \leq C_a|v|_{\alpha,\beta,M\times[0,T]}.$$

The notion of $C^{\alpha,\frac{\alpha}{2},\beta}[0,T]$, $C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$ and the other relevant norms will be clearified in Definition 2.2.

Remark 1.9. Theorem 1.8 is actually with respect to the real setting. Hopefully it can be applied to study real conical flows (such as Yamabe-flow) as well.

Remark 1.10. To prove the parabolic Schauder estimates, we follow Donaldson's idea in his proof of the elliptic Schauder estimates. Aside from Donaldson's work, Jefferes-Mazzeo-Rubinstein also proved a version of Schauder estimate for elliptic equations, by appealing to the "edge calculus machinery", while in this article we do everything by hand.

Theorem 1.8 depends on the following estimates on the heat kernel for the standard cone metric.

Theorem 1.11. *There exists a constant C as in Definition 2.6 with the following property. Suppose $\mathfrak{F}$ is a differential operator in $\mathfrak{P}$ (defined in (9)). Let $|\mathfrak{F}|$ be the order of the differential operator $\mathfrak{F}$. Then the following estimates on the heat kernel $H(x,y,t)$ of the standard cone metric $g_{E,\beta}$ (see Definition 6) hold.*

1. *Suppose $|\mathfrak{F}| = 1$. Suppose $t = 1$, $x \in A_{\frac{1}{\sqrt{s}}}$, $y \in A_{\frac{10}{\sqrt{s}}}$, $0 < s \leq \frac{1}{10000}$, then*
$$|\nabla_x H(x,y,1)| \leq Ce^{-\frac{1}{s}}.$$

2. *If $|\mathfrak{F}| \leq 2$, then for any x, y, we have*
$$\int_0^\infty |\mathfrak{F}H(x,y,\tau)|d\tau \leq C|x-y|^{-(m+|\mathfrak{F}|)}.$$

3. *Suppose $P(x) \geq \frac{|x-y|}{100^{100}}$. If $|\mathfrak{F}| = 3$, then we have*
$$\int_0^\infty |\mathfrak{F}H(x,y,\tau)|d\tau \leq C|y-x|^{-(m+3)}.$$

4. *For any x, y, we have*
$$\int_0^\infty |\nabla_x \frac{\partial}{\partial t} H(x,y,\tau)|d\tau \leq C|y-x|^{-(m+3)}.$$

5. *If $|\mathfrak{F}| \leq 4$, for any x, y and $0 < \alpha \leq 1$, we have*
$$\int_{\mathbb{R}^2\times\mathbb{R}^m} |\sup_{1\leq t\leq 2} \mathfrak{F}H(x,y,t)||x-y|^\alpha dy \leq C.$$

6. *For every second order spatial derivative operator $\mathfrak{D} \in \mathfrak{T}$, suppose $\rho = \min(\frac{1}{\beta}-1,\ 1)$, $|y| = 1$, and $|u_1|$, $|u_2| < \frac{1}{8}$, we have*
$$\int_0^\infty |\mathfrak{D}H(u_1,y,\tau) - \mathfrak{D}H(u_2,y,\tau)|d\tau \leq C|u_1-u_2|^\rho.$$

7. *Suppose $\rho = \min(\frac{1}{\beta}-1,\ 1)$, $|y| = 1$, and $|u_1|$, $|u_2| < \frac{1}{8}$, we have*
$$\int_0^\infty |\nabla_x H(u_1,y,\tau) - \nabla_x H(u_2,y,\tau)|d\tau \leq C|u_1-u_2|^\rho.$$

Proof. of Theorem 1.11: It's exactly a summarization of Lemma 7.1, 7.2, 7.3, 7.6, 7.5, Lemma 8.2, Lemma 9.1, and Proposition 9.2. ☐

Here the Properties (1-5) are satisfied by the heat kernel of the Euclidean space while the proofs are quite subtle. Property 6 and 7 are the only two properties which involve the exponent β. The difficulty of the proof is that the conical heat kernel is far less explicit than the Euclidean heat kernel or heat kernel in a smooth manifold. Thus, we are forced to deal with the representation formulas given in proposition 4.1 and Theorem 4.4. These representation formulas are due to Carslaw; and we reformulate them to apply to our case. In particular, the piecewise continuous (with respect to angle) representation formula in Theorem 4.4 is crucial. These representation formulas involve Bessel functions of the second kind, and some complicated integration of trignometric functions, and hyperbolic functions over angles.

Our proof is inspired by Donaldson's work [17]. There are some important difference in the technical level since we have to deal with a more complex situations. Restricted to the elliptic case, the function space we consider is exactly the same as Donaldson's $C^{2,\alpha,\beta}$ space in [17]. Our heat kernel estimates, restricted to the elliptic case, contain a little bit more information on the Green's function than the estimates in [17] (the Green function estimates in [17] is sufficient for elliptic Schauder estimates). Our proof also gives another approach to prove the estimates in [17] (see Section 7). The technical difference between our work and Donaldson's work on the Green's function in [17] is the following. In [17], Donaldson works from the heat kernel formula which involves Bessel functions of the first kind and uses a clever change varible in his proof. In our parabolic case, we employ two representation formulas of the heat kernel which are different from the one applied by Donaldson. One involves Bessel functions of the second kind and Weber's formula, the other one involves a contour integral which is essentially due to Carslaw. These different representation formulas are necessary (especially Carslaw's formula) in our parabolic case. Since we also need to understand the Hölder estimates over the time direction, we have to prove more global and more explicit properties of the heat kernel. While Donaldson shows us how to obtain Schauder estimates by working on Bessel functions, these new "additions" present difficult challenge for the proof our estimates. In summary, while ideas behind the heavy calculation are important, the sheer power of tensor manipulations and analysis of these Bessel functions are crucially needed.

For related work on the heat kernel of spaces with more genenral conical singularities, we refer the interested readers to the work by Mooers in [31] and by Mollers in [30].

The next diagram (Figure 1) is to indicate the relations of the lemmas, corollaries, theorems, and propositions in this article. Actually the whole article definitely can be summarized by Figure 1, aside from a few minor entanglements.



Figure 1: L means lemma, P means proposition, T means theorem, Cor means corollary. Arrows means imply.

Organization of this article: Most of the organization is illustrated in Figure 1. With respect to topics, this article can be divided into three parts:

1. proof of Theorem 1.2 assuming Theorem 1.8;
2. proof of Theorem 1.8 assuming Theorem 1.11;
3. proof of Theorem 1.11.

After defining the relevant norms explicitly and stating several more general theorems in Section 2, we carry out part 1 in section 3; part 2 in Sections 4, 5, 11. Both part 1 and part 2 are more or less standard, except some points where we should adapt polar coordinates into our framework. Part 3 is the most crucial part of this paper and it is also the most "non-regular" part. This is carried out in Section 6, 7, 8, 9, 10.

Acknowledgements: The second author would like to thank Professor Simon Donaldson for conversation on the theory of Bessel functions and his interest in this work. The second author is deeply grateful to Professor Xianzhe Dai, Professor Guofang Wei, and Prof Rugang Ye for offering and sponsoring the second author's first math job, for their interest in this work, for their continuous support and encouragements. Both authors would like to thank Haozhao Li, Weiyong He, Song Sun, and Kai Zheng very much for carefully reading the earlier versions of this paper.

2 Setting up of the main problems and more general statements.

First we would like to mention that, though some operators are not defined in the singular set, we still consider the domain of the functions to be the whole manifold M (instead of $M \setminus D$). The reason is that the functions and currents in our consideration are global, unless otherwise stated.

Let $m = 2n$ be a nonnegative integer. The local geometry of the conical metrics near the singularity hypersurfaces are like the following. Let $g_{E,\beta}$ be the standard cone metric on $\mathbb{R}^2 \times \mathbb{R}^m$, i.e.,

$$g_{E,\beta} = dr^2 + \beta^2 r^2 d\theta^2 + g_{\mathbb{R}^m}. \tag{6}$$

Here m is an even integer. In the subsequent work we will mainly use $\mathbb{R}^m$ instead of C^n, as our Schauder estimates are with respect to the real setting.

Now following Donaldson, let $z_j = s_j + \sqrt{-1}s_{\bar{j}}, j = 1, 2 \cdots n$ ($\bar{j} = j + n$). We consider a basis of $(1, 0)$ vectors as

$$\mathfrak{a} = \frac{1}{\sqrt{2}}(\frac{\partial}{\partial r} - \frac{\sqrt{-1}}{\beta r}\frac{\partial}{\partial \theta}), \frac{\partial}{\partial z_j}, j = 1 ... \frac{m}{2}. \tag{7}$$

Set $\xi = z^\beta = re^{i\beta\theta}$, notice that

$$\frac{\partial^2}{\partial \xi \partial \bar{\xi}} = \frac{1}{4}\Delta_\beta = \frac{1}{4}[\frac{\partial^2}{\partial r^2} + r^{-1}\frac{\partial}{\partial r} + \frac{1}{\beta^2}r^{-2}\frac{\partial^2}{\partial \theta^2}].$$

One can write the Laplacian of $g_{E,\beta}$ as

$$\Delta_{E,\beta} = \Delta_\beta + \Sigma_{i=1}^m \frac{\partial^2}{\partial s_i^2}.$$

When there is no confusion, we will denote $\Delta_{E,\beta}$ simply as Δ. Moreover, $\sqrt{-1}\partial\bar{\partial}$ operator is well defined under the basis

$$\frac{\partial^2}{\partial \xi \partial \bar{\xi}}, \mathfrak{a}\frac{\partial}{\partial \bar{z}_i}, \bar{\mathfrak{a}}\frac{\partial}{\partial z_i}, \frac{\partial^2}{\partial z_i \partial \bar{z}_j}, 1 \leq i, j \leq n.$$

To state the main properties of the heat kernel, we need to introduce some linear, differential operators (up to 4th orders). First, we write ∇_x as the gradient operator in $\mathbb{R}^2 \times \mathbb{R}^m$ with respect to a natural basis

$$\frac{\partial}{\partial r}, \frac{1}{\beta r}\frac{\partial}{\partial \theta}, \frac{\partial}{\partial s_i}(1 \leq i \leq m).$$

Secondly, we can introduce a set of real, second order operators $\mathfrak{T}$ as:

$$\mathfrak{T} = \{\frac{\partial^2}{\partial r \partial s_i},\ i \leq m;\ \frac{\partial^2}{\partial s_i \partial s_j},\ i, j \leq m;\ \frac{1}{r}\frac{\partial^2}{\partial s_i \partial \theta},\ i \leq m\}. \tag{8}$$

Finally, the set of operators $\mathfrak{P}$ we need to study in this paper as

$$\mathfrak{P} = \{\nabla_x, \frac{\partial}{\partial t}, \mathfrak{D}, \nabla_x\nabla_y; \nabla_x\mathfrak{D}, \nabla_x\frac{\partial}{\partial t}; \frac{\partial}{\partial t}\mathfrak{D}, \frac{\partial^2}{\partial t^2}|\mathfrak{D} \in \mathfrak{T}\}. \tag{9}$$

With the $\sqrt{-1}\partial\bar{\partial}$ operator in mind, we define the Sobolev space $W^{2,2}(M \times [0, T])$ by the following norm.

$$\begin{aligned} & |\widehat{u}|_{W^{2,2}(M\times[0,T])} \qquad (10) \\ \triangleq\ & |\frac{\partial \widehat{u}}{\partial t}|_{L^2(M\times[0,T])} + |\sqrt{-1}\partial\bar{\partial}\widehat{u}|_{L^2(M\times[0,T])} + |\nabla\widehat{u}|^{(1)}_{L^2(M\times[0,T])} + |\widehat{u}|_{L^2(M\times[0,T])}. \end{aligned}$$

This Sobolev norm has a good application in Proposition 5.8.

Let Ω be a domain in $\mathbb{R}^2 \times \mathbb{R}^m$. For any point $x \in \Omega$ we denote d_x to be the distance from x to the boundary $\partial\Omega$. Notice that the points in the singular set $\{0\} \times R^m \cap \ int(\Omega)$ are never considered to be boundary points, and this is quite crucial in this article. For any two points $x, y \in \Omega$, we set

$$d_{x,y} = \min(d_x, d_y),\ d_x \text{ is the spatial distance from } x \text{ to } \partial\Omega.$$

Similarly, we set

$$l_{(x,t_1),(y,t_2)} = \min(l_{(x,t_1)}, l_{(y,t_2)}),$$

where the $l_{(x,t)}$ is defined as

$$l_{(x,t)} = \min\{d_x,\ |t - T_1|^{\frac{1}{2}}\}.$$

For any $[T_1, T_2] \subset \mathbb{R}^+$ and for any function $\widehat{u}$ over $\Omega \times [T_1, T_2]$, we introduce the usual semi norms as

$$\begin{aligned}
[\widehat{u}]_{0,\Omega\times[T_1,T_2]} &= \sup_{\Omega\times[T_1,T_2]} |\widehat{u}(x,t)|, \\
[\widehat{u}]_{\alpha,\Omega\times[T_1,T_2]} &= \sup_{(x,t),(y,t)\in\Omega\times[T_1,T_2]} \frac{|\widehat{u}(x,t) - \widehat{u}(y,t)|}{|x-y|^\alpha}, \\
[\widehat{u}]_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]} &= \sup_{(x,t_1),(y,t_2)\in\Omega\times[T_1,T_2]} \frac{|\widehat{u}(x,t_1) - \widehat{u}(y,t_2)|}{|x-y|^\alpha + |t_1 - t_2|^{\frac{\alpha}{2}}}.
\end{aligned}$$

Now, we define weighted Schauder norms (with spatial distance as weights) as:

$$\begin{aligned}
[\widehat{u}]^{(k)}_{0,\Omega\times[T_1,T_2]} &= \sup_{\Omega\times[T_1,T_2]} d_x^k |\widehat{u}(x,t)|, \\
[\widehat{u}]^{(k)}_{\alpha,\Omega\times[T_1,T_2]} &= \sup_{(x,t),(y,t)\in\Omega\times[T_1,T_2]} d_{x,y}^{k+\alpha} \frac{|\widehat{u}(x,t) - \widehat{u}(y,t)|}{|x-y|^\alpha}, \\
[\widehat{u}]^{(k)}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]} &= \sup_{(x,t_1),(y,t_2)\in\Omega\times[T_1,T_2]} d_{x,y}^{k+\alpha} \frac{|\widehat{u}(x,t_1) - \widehat{u}(y,t_2)|}{|x-y|^\alpha + |t_1 - t_2|^{\frac{\alpha}{2}}}.
\end{aligned}$$

We also define parabolic weighted Schauder norms (with parabolic distance as weights) as:

$$\begin{aligned}
[\widehat{u}]^{[k]}_{0,\Omega\times[T_1,T_2]} &= \sup_{\Omega\times[T_1,T_2]} l_{x,t}^k |\widehat{u}(x,t)|, \\
[\widehat{u}]^{[k]}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]} &= \sup_{(x,t_1),(y,t_2)\in\Omega\times[T_1,T_2]} l_{(x,t_1);(y,t_2)}^{k+\alpha} \frac{|\widehat{u}(x,t_1) - \widehat{u}(y,t_2)|}{|x-y|^\alpha + |t_1 - t_2|^{\frac{\alpha}{2}}}.
\end{aligned}$$

We define the norm $|\cdot|^{*}_{2,\alpha,\Omega\times[T_1,T_2]}$ as

$$\begin{aligned}
&|\widehat{u}|^{*}_{2,\alpha,\Omega\times[T_1,T_2]}\\
\triangleq\;& [\frac{\partial\widehat{u}}{\partial t}]^{(2)}_{\alpha,\Omega\times[T_1,T_2]}+[i\partial\bar{\partial}\widehat{u}]^{(2)}_{\alpha,\Omega\times[T_1,T_2]}+|i\partial\bar{\partial}\widehat{u}|^{(2)}_{0,\Omega\times[T_1,T_2]}\\
&+|\frac{\partial\widehat{u}}{\partial t}|^{(2)}_{0,\Omega\times[T_1,T_2]}+|\nabla\widehat{u}|^{(1)}_{0,\Omega\times[T_1,T_2]}+|\widehat{u}|_{0,\Omega\times[T_1,T_2]}.
\end{aligned}$$

Finally, we define the space $\widehat{C}^{2+\alpha,1+\frac{\alpha}{2},\beta}_{\star}\{\Omega\times[T_1,T_2]\}$ (with spatial distance as weights) by the norm

$$\begin{aligned}
&|\widehat{u}|^{*}_{2+\alpha,1+\frac{\alpha}{2},\Omega\times[T_1,T_2]}\\
\triangleq\;& [i\partial\bar{\partial}\widehat{u}]^{(2)}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]}+[\frac{\partial\widehat{u}}{\partial t}]^{(2)}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]}\\
&+[\frac{\partial\widehat{u}}{\partial t}]^{(2)}_{0,\Omega\times[T_1,T_2]}+[i\partial\bar{\partial}\widehat{u}]^{(2)}_{0,\Omega\times[T_1,T_2]}+|\nabla\widehat{u}|^{(1)}_{0,\Omega\times[T_1,T_2]}+|\widehat{u}|_{0,\Omega\times[T_1,T_2]};
\end{aligned}$$

and the space $\widehat{C}^{2+\alpha,1+\frac{\alpha}{2},\beta}_{[\star]}\{\Omega\times[T_1,T_2]\}$ (with parabolic distance as weights) by the norm

$$\begin{aligned}
&|\widehat{u}|^{[*]}_{2+\alpha,1+\frac{\alpha}{2},\Omega\times[T_1,T_2]}\\
\triangleq\;& [i\partial\bar{\partial}\widehat{u}]^{[2]}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]}+[\frac{\partial\widehat{u}}{\partial t}]^{[2]}_{\alpha,\frac{\alpha}{2},\Omega\times[T_1,T_2]}\\
&+[\frac{\partial\widehat{u}}{\partial t}]^{[2]}_{0,\Omega\times[T_1,T_2]}+[i\partial\bar{\partial}\widehat{u}]^{[2]}_{0,\Omega\times[T_1,T_2]}+|\nabla\widehat{u}|^{[1]}_{0,\Omega\times[T_1,T_2]}+|\widehat{u}|_{0,\Omega\times[T_1,T_2]}.
\end{aligned}$$

Actually in most of the cases we will consider the polydisks centered at points in the divisor. Namely we will frequently appeal to the following definition.

Definition 2.1. Given p in the singular set $(0)\times\mathbb{R}^m$, we consider $A_R(p)$ as

$$A_R(p)=D_R(p)\times\widehat{B}_R(p),$$

where $D_R(p)$ is the disk of radius R in the $\mathbb{R}^2$ component and $\widehat{B}_R(p)$ is the ball of radius R in the $\mathbb{R}^m$ component. When the center is 0 we abbreviate $A_R(0)$ as A_R. The advantage of using such domains can be seen, for example, when we estimate Z_6 in the proof of proposition 5.4.

We wish to point out that, unlike in the paper [17], we allow the hypersurface D is the disjoint union of smooth irreducible divisors. Set

$$\Lambda=\Sigma^{l}_{j=1}2\pi(1-\beta_i)D_i,$$

where each D_i is a connected divisor. For any ϵ small, denote the tubular neighborhood $T_\epsilon(D)$ with radius ϵ of the hypersurface D. Set $\nu > 0$ to be small enough so that the following decomposition holds

$$T_{2\nu}(D) \subset \bigcup_{i=1}^{l} \bigcup_{k=1}^{l_{D_i}} N_k(D_i)$$

(such that in each $N_k(D_i)$, D_i is the zero locus of some holomorphic function z_0 in $N_k(D_i)$. Suppose ι_{β_i} is the map defined by

$$\begin{array}{llll} \iota_{\beta_i} : N_k(D_i) & \to & \mathbb{R}^2 \times \mathbb{C}^n \\ \qquad (z_0, z_1 ......z_n) & \to & (|z_0|^{\beta_i - 1} z_0, z_1, ...z_n). \end{array}$$

Using this map, we define

$$|u|_{2,\alpha,\Lambda,M\times[T_1,T_2]} = \Sigma_{i=1}^{l} \Sigma_{k=1}^{l_{D_i}} |\widehat{u}|^*_{2,\alpha,\iota_{\beta_i}[N_k(D_i)]\times[T_1,T_2]} + |u|^*_{2,\alpha,[M\setminus T_\nu(D)]\times[T_1,T_2]} \quad (11)$$

and

$$\begin{aligned} & |u|_{2+\alpha,\ 1+\frac{\alpha}{2},\Lambda,M\times[T_1,T_2]} \\ = \ & \Sigma_{i=1}^{l} \Sigma_{k=1}^{l_{D_i}} |\widehat{u}|^*_{2+\alpha,\ 1+\frac{\alpha}{2},\iota[N_k(D)]\times[T_1,T_2]]} + |u|^*_{2+\alpha,\ 1+\frac{\alpha}{2},[M\setminus T_\nu(D)]\times[T_1,T_2]}. \quad (12) \end{aligned}$$

Definition 2.2. We define the space $C^{2,\alpha,\Lambda}[T_1, T_2]$ by the norm $|\cdot|_{2,\alpha,\Lambda,M\times[T_1,T_2]}$ and the space $C^{2+\alpha,1+\frac{\alpha}{2},\Lambda}[T_1, T_2]$ by $|\ \cdot\ |_{2+\alpha,\ 1+\frac{\alpha}{2},\Lambda,M\times[T_1,T_2]}$.

When $\beta_i = \beta$ for every i, then we simply replace Λ by β to denote the space and norms

$$C^{2+\alpha,1+\frac{\alpha}{2},\Lambda}[T_1, T_2]\ (C^{2,\alpha,\Lambda}[T_1, T_2]),\ |\ \cdot\ |_{2+\alpha,1+\frac{\alpha}{2},\Lambda,M\times[T_1,T_2]}\ (|\ \cdot\ |_{2,\alpha,\Lambda,M\times[T_1,T_2]}),$$

as

$$C^{2+\alpha,1+\frac{\alpha}{2},\beta}[T_1, T_2]\ (C^{2,\alpha,\beta}[T_1, T_2]),\ |\ \cdot\ |_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[T_1,T_2]}\ (|\ \cdot\ |_{2,\alpha,\beta,M\times[T_1,T_2]}).$$

The other norms concerning Λ should have Λ replaced by β also.

A time-independent function u is said to be in $C^{2,\alpha,\beta}$ ($C^{2,\alpha,\Lambda}$), if u is in $C^{2,\alpha,\beta}[T_1, T_2]$ ($C^{2,\alpha,\Lambda}[T_1, T_2]$. Since u does not depend on time, the definition does not depend on T_1, T_2 at all.

Remark 2.3. Similar to norms in (11) and (12), we can use the transformations ι_{β_i} to define weaker norms

$$|\ \cdot\ |_{1,\alpha,\Lambda,M\times[T_1,T_2]};\quad |\ \cdot\ |_{1+\alpha,\frac{1}{2}+\frac{\alpha}{2},\Lambda,M\times[T_1,T_2]}$$

and

$$|\ \cdot\ |_{\alpha,\Lambda,M\times[T_1,T_2]};\quad |\ \cdot\ |_{\alpha,\frac{\alpha}{2},\Lambda,M\times[T_1,T_2]}.$$

Remark 2.4. When u is a time independent function over M, we may define the norm $\mid u \mid_{2,\alpha,\Lambda,M}$ as in [17]. To be more precise, the norm $\mid u \mid_{2,\alpha,\Lambda,M}$ is also equivalent to $\mid u \mid_{2,\alpha,\Lambda,M\times[T_1,T_2]}$ if we view u to be a space time function though u doesn't depend on time.

Definition 2.5. A time-dependent family $\{a(t),\ t\in[0,T]\}$ of (α,β) metrics is said to be $C^{\alpha,\frac{\alpha}{2},\beta}[0,T]$ if there exist a $\phi(t)\in C^{2+\alpha,1+\frac{\alpha}{2},\beta}(M\times[0,T])$ and a smooth metric $\acute{\omega}$ such that

$$a(t)=\acute{\omega}+i\partial\bar{\partial}\phi(t).$$

In the general case, we should consider the reference metric ω_0 as[1]

$$\omega_\Lambda=\omega^{'}+\delta i\partial\bar{\partial}[|S_1|_1^{2\beta_1}\times|S_2|_2^{2\beta_2}\times......|S_i|_i^{2\beta_i}......|S_l|_l^{2\beta_l}].$$

and define the (α,Λ) metrics as in definition 1.1 (with β replaced by Λ). Then, more general theorems hold (comparing with Theorems 1.2, 1.7, 1.8).

Definition 2.6. Since we are dealing with many estimates in this article, it's necessary to illustrate a principle of the dependency of C in every theorem, proposition, corollary and lemma. Namely, we make the following important convention in this article:

Without further notice, the "C" in each estimate means a constant depending on the dimension $m+2$, the angle β, the α (and $\acute{\alpha}$ if any) in the same estimate or in the corresponding theorem (proposition, corollary, lemma). We add subindex to the "C" if it depends on more factors than the above three. Moreover, the "C" in different places might be different.

Theorem 2.7. *Suppose g_0 is an $(\acute{\alpha},\Lambda)$ conical Kähler metric in $(M,\Sigma_{j=1}^{l}(1-\beta_j)D_j)$ where $\alpha'\in(0,\min\{\min_{1\le j\le l}[\frac{1}{\beta_j}-1],1\})$. For any $\alpha\in(0,\alpha')$, and constant number μ, there exists an $T_0>0$ (which depends on g_0) such that the conical Kähler-Ricci flow equation*

$$\frac{\partial\omega_g}{\partial t}=\beta\omega_g-Ric(g)+2\pi\Sigma_{j=1}^{l}(1-\beta_j)D_j,\ \omega_g=\omega_{g_0}\ \textit{when}\ t=0$$

admits a solution $g(t)$, $t\in[0,T_0]$ which is smooth (both in space and time) in $(M\setminus D)\times[0,T_0]$. Moreover we have

- *for every $t\in[0,T_0]$, $g(t)$ is an (α,Λ) conical metric in $(M,\Sigma_{j=1}^{l}(1-\beta_j)D_j)$;*

---

[1] Note that δ must be small enough.

- *$g(t)$ is a $C^{\alpha,\frac{\alpha}{2},\Lambda}[0,T_0]$-family of conical metrics (see Definition 2.5).*
- *for any $0<\widehat{\alpha}\leq\alpha$, $g(t)$ is the unique solution of (2) in the class of $C^{\widehat{\alpha},\frac{\widehat{\alpha}}{2},\Lambda}[0,T]$-family of metrics.*

Theorem 2.8. *Suppose $0<T<\infty$ and $\{a(t),\ t\in[0,T]\}$ is a $C^{\alpha,\frac{\alpha}{2},\Lambda}[0,T]$-family of (α,Λ) metrics (See Definition 2.5). Then there exists a constant C_a (depending on $a(t)$) with the following property. For any $v\in C^{\alpha,\frac{\alpha}{2},\Lambda}[0,T]$, the parabolic equation*

$$\frac{\partial u}{\partial t}=\Delta_{a(t)}u+v,\ \ u(0,x)=0 \tag{13}$$

admits a unique solution u in $C^{2+\alpha,1+\frac{\alpha}{2},\Lambda}[0,T]$. Moreover, the following estimates hold.

$$|u|_{2+\alpha,1+\frac{\alpha}{2},\Lambda,M\times[0,T]}\leq C_a|v|_{\alpha,\frac{\alpha}{2},\Lambda,M\times[0,T]};$$
$$|u|_{2,\alpha,\Lambda,M\times[0,T]}\leq C_a|v|_{\alpha,\Lambda,M\times[0,T]}.$$

3 Proof of Theorem 1.2 and Theorem 1.7 assuming Theorem 1.8.

In this section we assume Theorem 1.8 to be right and prove the short time existence of CKRF. The proof is actually regular and no special conical structure is involved. For the sake of being rigorous we shall include the proof here. Since Theorem 1.8 is an independent part, we left it to later discussions.

Proof. of Theorem 1.7: Actually Theorem 1.7 is a direct conclusion of Theorem 1.8 and the iteration argument. Since we lack an exact reference on the result we want, we include the crucial detail here. We still denote $C_0^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$ as the subspace of functions with zero initial value of $C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$. For brevity, we only prove in detail the short time existence result assuming $\mu=0$. The result for all μ follows from exactly the same method.

We consider the following two equations

$$\frac{\partial\phi}{\partial t}=\log\frac{(\omega_0+\sqrt{-1}\partial\bar{\partial}\phi)^n}{\omega_0^n}+f;\phi=0\text{ when }t=0. \tag{14}$$

$$\frac{\partial u}{\partial t}=\Delta_{\omega_0}u+f,\ \ u=0\text{ when }t=0. \tag{15}$$

Substracting (15) from (14) we end up with the followin equation.

$$\frac{\partial v}{\partial t} = \log \frac{(\omega_u + \sqrt{-1}\partial\bar{\partial}v)^n}{\omega_u^n} + F(u); v = 0 \text{ when } t = 0. \tag{16}$$

In the above equation, we define that

$$\omega_u = (\omega_0 + \sqrt{-1}\partial\bar{\partial}u)^n,\ v = \phi - u,\ F(u) = \log \frac{\omega_u^n}{\omega_0^n} - \Delta_{\omega_0} u.$$

Notice that when T is small enough, ω_u is still a uniformly elliptic family of metric. A crucial advantage of this trick of substraction is that

$$F(u) = 0 \text{ when } t = 0,$$

which implies that the linearized operator is a contraction for short time.

To continue, we define $L = \frac{\partial}{\partial t} - \Delta_{\omega_u} - F(u)$. Then this defines an operator

$$L : C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T] \to C^{\alpha, \frac{\alpha}{2}, \beta}[0, T].$$

By Theorem 1.8, this is an invertible operator with $|L^{-1}|$ uniformly bounded. Set

$$E(v) = \log \frac{(\omega_u + i\partial\bar{\partial}v)^n}{\omega_u^n} - \Delta_{\omega_u}.$$

Consider the operator

$$\Phi = L^{-1} \circ E : C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T] \to C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T].$$

We shall show in the following that Φ is a contracting map in a small ball B_{ϵ_1} (centered at the zero function) in $C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T]$. For any $v_1, v_2 \in C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T]$, there exist $u_1, u_2 \in C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T]$ such that

$$Lu_1 = E(v_1) \qquad \text{and } \ Lu_2 = E(v_2)$$

or equivalently

$$\Phi(v_1) = u_1 \qquad \text{and } \ \Phi(v_2) = u_2.$$

Thus we obtain the equation

$$L(u_1 - u_2) = E(v_1) - E(v_2).$$

Note that E is a quadratic operator. Suppose $|v_1|, |v_2| \le \epsilon_2$ and ϵ_2 is small enough, we have that

$$|E(v_1) - E(v_2)|_{\alpha, \frac{\alpha}{2}, \beta, M\times[0,T]} \le \epsilon_3 |v_1 - v_2|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]}$$

By Theorem 1.8, we have

$$\begin{aligned} |u_1 - u_2|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} &\leq C_1|E(v_1) - E(v_2)|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \\ &\leq C_1 \cdot \epsilon_3 \cdot |v_1 - v_2|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]}. \end{aligned}$$

Then making ϵ_2 sufficiently small, ϵ_3 can be small enough too. Thus, we have:

$$\begin{aligned} & |\Phi(v_1) - \Phi(v_2)|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \\ = \; & |u_1 - u_2|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \leq \frac{1}{2}|v_1 - v_2|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]}. \end{aligned}$$

Then, Φ is a contraction map in B_{ϵ_2}. To show that there is a fixed point, it suffices to show the first iteration is small (provided the time is short). We have the following crucial lemma about the first iteration.

Claim 3.1. *Let $v_0 \equiv 0$. For any $\epsilon_4 > 0$, there exists a T small enough such that*

$$|\Phi(v_0)|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \leq \epsilon_4.$$

To prove claim 3.1, it suffices to use the Schauder estimate over time. This is achieved as follows. By Theorem 1.8 we know that $F(u) \in C^{\acute{\alpha},\frac{\acute{\alpha}}{2},\beta}[0,T]$. Notice by our choice we have that $\alpha < \acute{\alpha}$. Then using the crucial fact that $F(u) = 0$ when $t = 0$, we trivially get

$$|F(u)|_{0,M\times[0,T]} \leq T^{\frac{\acute{\alpha}}{2}}[F(u)]_{\acute{\alpha},\frac{\acute{\alpha}}{2},\beta,M\times[0,T]}.$$

Denote $\Phi(v_0)$ as v_1, notice that

$$\frac{\partial v_1}{\partial t} - \Delta_{\omega_u} v_1 - F(u) = 0.$$

Then by applying Theorem 1.8, we have when T is small enough that

$$\begin{aligned} & |v_1|_{2+\alpha,1+\frac{\alpha}{2},\beta,M\times[0,T]} \\ \leq \; & C|F(u)|_{\alpha,\frac{\alpha}{2},\beta,M\times[0,T]} \\ = \; & C\{|F(u)|_{0,M\times[0,T]} + [F(u)]_{\alpha,\frac{\alpha}{2},\beta,M\times[0,T]}\} \\ \leq \; & CT^{\frac{\acute{\alpha}}{2}}[F(u)]_{\acute{\alpha},\frac{\acute{\alpha}}{2},\beta,M\times[0,T]} + CT^{\frac{\acute{\alpha}-\alpha}{2}}|F(u)|_{\acute{\alpha},\frac{\acute{\alpha}}{2},\beta,M\times[0,T]} \leq \epsilon_4. \end{aligned}$$

Then claim 3.1 is proved.

Now we are ready to define the iteration process. Set $v_0 = 0$ and inductively, we define

$$v_{k+1} = \Phi(v_k), \qquad \text{where } \ k = 0, 1, 2, \cdots .$$

Then combining claim 3.1 we have for T small that $|v_k|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]} \leq \epsilon_3$ (then Φ is a contraction over the sequence). Hence

$$\begin{array}{rcl} |v_{k+1} - v_k|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]} & = & |\Phi(v_k) - \Phi(v_{k-1})|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]} \\ & \leq & \frac{1}{2}|v_k - v_{k-1}|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]} \\ & \leq & \frac{1}{2^k}|v_1 - v_0|_{2+\alpha, 1+\frac{\alpha}{2}, \beta, M\times[0,T]} \leq \frac{\epsilon_4}{2^k}. \end{array}$$

Then it is easy to see that the sequence $\{v_k\}$ is a Cauchy sequence in $C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T]$. Thus, $\{v_k\}$ converge to a fixed point $v_\infty \in C_0^{2+\alpha, 1+\frac{\alpha}{2}, \beta}[0, T]$ such that

$$\Phi(v_\infty) = v_\infty.$$

In other words,

$$\frac{\partial v_\infty}{\partial t} = \log \frac{(\omega_u + i\partial\bar{\partial} v_\infty)^n}{\omega_u^n} + F(u).$$

Therefore $\phi = v_\infty - u$ is a solution to (14).

□

Proof. of Theorems 1.2, 2.7: Again we only say a few words on Theorem 1.2. Theorem 2.7 is similar.

Now, recall

$$f = h_{\omega_D} + \log \frac{\omega_0^n}{\omega_D^n}.$$

Since ω is $(\acute{\alpha}, \beta)$, then using 4 we see that $f \in C^{\acute{\alpha}, \beta}$. Thus by applying Theorem 1.7, the proof is complete. □

4 Representation formulas for the heat kernel.

In this section, we give a representation formula for the heat Kernel with respect to the conical model metric $g_{E,\beta}$ over $\mathbb{R}^2 \times \mathbb{R}^m$.

As in [49], we denote $I_\mu(z)$ as the Bessel function of second kind with order μ and $J_\mu(z)$ as the Bessel function of first kind with order μ.

First we recall the Weber's formula

$$\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}I_\mu(\frac{rr^{'}}{2t})=\int_0^\infty e^{-\lambda^2t}J_\mu(\lambda r)J_\mu(\lambda r^{'})\lambda d\lambda, \tag{17}$$

First, for the sake of completeness we give a proof of the heat kernel formula when $m=0$. . We believe this proof is well known to experts.

Proposition 4.1. *The heat kernel of* $\{R^2,\ dr^2+\beta^2r^2d\theta^2\}$ *is*

$$\begin{aligned}
& \widehat{H}(r,\theta,r^{'},\theta^{'},t) \\
= & \frac{1}{2\pi\beta}\int_0^\infty e^{-\lambda^2t}J_0(\lambda r)J_0(\lambda r^{'})\lambda d\lambda \\
& +\frac{1}{2\pi\beta}\Sigma_{k\geq 1}2\cos(k(\theta-\theta^{'}))\int_0^\infty e^{-\lambda^2t}J_{\frac{k}{\beta}}(\lambda r)J_{\frac{k}{\beta}}(\lambda r^{'})\lambda d\lambda \\
= & \frac{1}{2\pi\beta}\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}I_0(\frac{rr^{'}}{2t})+\frac{1}{2\pi\beta}\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}\Sigma_{k\geq 1}2\cos(k(\theta-\theta^{'}))I_{\frac{k}{\beta}}(\frac{rr^{'}}{2t}).
\end{aligned}$$

Proof. of Proposition 4.1: First, we prove by direct computation that

$$(\Delta_\beta-\frac{\partial}{\partial t})\widehat{H}=0.$$

Secondly, we want to prove that

$$\lim_{t\to 0}\beta\int_{-\pi}^{\pi}\int_0^{+\infty}\widehat{H}(r,\theta,r^{'},\theta^{'},t)f(r^{'},\theta^{'})r^{'}dr^{'}d\theta^{'}=f(r,\theta).$$

We claim that the following holds.

Claim 4.2. *Suppose* $v>-\frac{1}{2}$ *and* $r>0$, *we have that*

$$\lim_{t\to 0}\int_0^{+\infty}\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}I_v(\frac{rr^{'}}{2t})f(r^{'},\theta^{'})r^{'}dr^{'}=f(r,\theta^{'}).$$

Proof. Suppose $r>0$, set $u=\frac{r^{'}-r}{2\sqrt{t}}$. We have

$$\begin{aligned}
& \lim_{t\to 0}\int_0^{+\infty}\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}I_v(\frac{rr^{'}}{2t})f(r^{'},\theta^{'})r^{'}dr^{'} \\
= & \lim_{t\to 0}\int_{-\infty}^{+\infty}\frac{1}{\sqrt{t}}e^{-u^2}I_v(\frac{r^2+2ru\sqrt{t}}{2t})e^{-\frac{r^2+2ru\sqrt{t}}{2t}}f(r+2u\sqrt{t},\theta^{'})(r+2u\sqrt{t})du
\end{aligned}$$

According to formula (1) of $I_v(x)$ in 7.25 of [49], we have

$$\begin{aligned} & \lim_{t\to 0}\frac{1}{\sqrt{t}}I_v(\frac{r^2+2ru\sqrt{t}}{2t})e^{-\frac{r^2+2ru\sqrt{t}}{2t}}(r+2u\sqrt{t}) \\ = & \lim_{t\to 0}\frac{1}{\sqrt{\pi}\sqrt{\frac{r^2+2ru\sqrt{t}}{t}}} * \frac{\int_0^{\frac{r^2+2ru\sqrt{t}}{t}} w^{v-\frac{1}{2}}e^{-w}dw}{\Gamma(v+\frac{1}{2})} * \frac{r+2u\sqrt{t}}{\sqrt{t}} \\ = & \frac{1}{\sqrt{\pi}}. \end{aligned}$$

It follows that

$$\begin{aligned} & \lim_{t\to 0}\int_0^{+\infty}\frac{1}{2t}e^{-\frac{r^2+r^{'2}}{4t}}I_v(\frac{rr^{'}}{2t})f(r^{'},\theta^{'})r^{'}dr^{'} \\ = & \frac{1}{\sqrt{\pi}}\int_{-\infty}^{+\infty}e^{-u^2}duf(r,\theta^{'}) = f(r,\theta^{'}). \end{aligned}$$

Then claim 4.2 is proved. □

Now we are ready to prove proposition 4.1. From Fourier Series theory we trivially have

$$\begin{aligned} & \lim_{t\to 0}\beta\int_{-\pi}^{\pi}\int_0^{+\infty}H(r,r^{'},\theta,\theta^{'},t)f(r^{'},\theta^{'})r^{'}dr^{'}d\theta^{'} \\ = & \frac{1}{2\pi}\int_{-\pi}^{\pi}f(r,\theta^{'})d\theta^{'}+\frac{1}{2\pi}\Sigma_{k\geq 1}2\int_{-\pi}^{\pi}f(r,\theta^{'})\cos(k(\theta-\theta^{'}))d\theta^{'} \\ = & f(r,\theta). \end{aligned}$$

□

Remark 4.3. One should notice that when $r=0$, the heat kernel does not tend to the $\delta-$function anymore when $t\to 0$. Nevertheless, this does not affect Lemma 4.5 to be true since we assume $r\neq 0$ in it.

Thus it's easy to get the expression for the heat kernel $H(x,y,t)$ of our model conical metric $g_{E,\beta}$:

$$\begin{aligned} H(x,y,t) = & H(r,\theta,r^{'},\theta^{'},R,t) \\ = & \frac{1}{\beta(4\pi t)^{\frac{m}{2}+1}}e^{-\frac{r^2+r^{'2}+R^2}{4t}}[I_0(\frac{rr^{'}}{2t})+\Sigma_{k\geq 1}2\cos(k(\theta-\theta^{'}))I_{\frac{k}{\beta}}(\frac{rr^{'}}{2t})] \\ = & \frac{2t}{\beta(4\pi t)^{\frac{m}{2}+1}}e^{-\frac{R^2}{4t}}[\int_0^{\infty}e^{-\lambda^2 t}J_0(\lambda r)J_0(\lambda r^{'})\lambda d\lambda \\ + & \Sigma_{k\geq 1}2\cos(k(\theta-\theta^{'}))\int_0^{\infty}e^{-\lambda^2 t}J_{\frac{k}{\beta}}(\lambda r)J_{\frac{k}{\beta}}(\lambda r^{'})\lambda d\lambda]. \end{aligned} \tag{18}$$

Here $x = (r, \theta, \widehat{x})$, $y = (r^{'}, \theta^{'}, \widehat{y})$. $R = |\widehat{x} - \widehat{y}|$ is the tangential distance parallel to $\{0\} \times R^m$.



Figure 2 for the contour $\mathfrak{A}$

Theorem 4.4. *(Reformulation of Carslaw's results in [7]) The heat kernel of the flat conical Kähler metric* $g_{E,\beta}$ *is*

$$H(r, \theta, \acute{r}, \theta^{'}, R, t) = \frac{1}{2\pi\beta} \frac{1}{(4\pi t)^{\frac{m}{2}+1}} e^{-\frac{r^2+\acute{r}^2+R^2}{4t}} P[\frac{r\acute{r}}{2t}, \beta(\theta^{'} - \theta)].$$

In the above formula, P *is defined as*

$$P(z, v) \quad \triangleq \quad \int_{\mathfrak{A}} e^{z\cos a} \frac{1}{1 - e^{-\frac{i(a-v)}{\beta}}} da.$$

P can also be expressed as

$$\begin{aligned} & P(z,v) \\ = \;& 2\pi[I_0(z)+\Sigma_{k\geq 1}2I_{\frac{k}{\beta}}(z)\cos\frac{kv}{\beta}]. \\ = \;& 2\pi\beta\Sigma_{k,\ -\pi<v+2k\beta\pi<\pi}e^{z\cos(v+2k\beta\pi)}+E(z,v) \\ & +\pi\beta\Sigma_{k,v+2k\beta\pi=\pm\pi}e^{-z}. \end{aligned} \tag{19}$$

Here

$$E(z,v)=\int_0^{\infty}e^{-z\cosh y}\frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]}dy,$$

and $\mathfrak{A}$ *is the path in Figure 2 above the theorem (which is the same as FIG 1 of [7] with argument translated by* θ*).*

Proof. of Theorem 4.4: The first expression of the heat kernel is exactly from the results in Section 2 iof [7]. The second expression in the first line of formula (19) (which concerns the Bessel function of the second kind I) is also exactly from [7]. Thus we only have to prove the expression in the second line of formula (19). Working from the formula

$$P(z,v)=\int_{\mathfrak{A}}e^{z\cos a}\frac{1}{1-e^{-\frac{i(a-v)}{\beta}}}da, \tag{20}$$

the singularities of this integrand are

$$a_k=v+2k\beta\pi,\ -\pi\leq v+2k\beta\pi\leq\pi.$$

We can of course deform $\mathfrak{A}$ continuously away from these singularities to get a integral over the L_1 and L_2 in Figure 2, together with the residue integrals around the singularities a_k. At those a_k ($a_k=v+2k\beta$) such that $-\pi<a_k<\pi$, the residues are $2\pi\beta$. At those a_k such that $a_k=v+2k\beta\pi=\pi$ or $-\pi$, the contribution of the half circle with radius tending to 0 is

$$\lim_{\epsilon\to 0}\int_A e^{z\cos a}\frac{1}{1-e^{-\frac{i(a-v)}{\beta}}}da=\pi\beta e^{z\cos(\pm\pi)}=\pi\beta e^{-z}.$$

To prove the formula for $E(z,v)$ in formula 19, it suffices to notice that over L_1, $a=-\pi+iy,\ y\in(-\infty,+\infty)$; over L_2, $a=\pi+iy,\ y\in(+\infty,-\infty)$. Thus,

$$\begin{aligned}
&\int_{L_1+L_2} e^{z[\cos a]}\frac{1}{1-e^{-\frac{i(a-v)}{\beta}}}da\\
= &\int_{-\infty}^{\infty} e^{-z\cosh y}\frac{i}{1-e^{\frac{iv}{\beta}}e^{\frac{y}{\beta}}e^{\frac{i\pi}{\beta}}}dy - \int_{-\infty}^{\infty} e^{-z\cosh y}\frac{i}{1-e^{\frac{iv}{\beta}}e^{\frac{y}{\beta}}e^{-\frac{i\pi}{\beta}}}dy\\
= &\int_0^{\infty} e^{-z\cosh y}F(v,y)dy,
\end{aligned}$$

where

$$F(v,y)=\frac{i}{1-e^{\frac{iv}{\beta}}e^{\frac{y}{\beta}}e^{\frac{i\pi}{\beta}}}+\frac{i}{1-e^{\frac{iv}{\beta}}e^{-\frac{y}{\beta}}e^{\frac{i\pi}{\beta}}}-\frac{i}{1-e^{\frac{iv}{\beta}}e^{\frac{y}{\beta}}e^{-\frac{i\pi}{\beta}}}-\frac{i}{1-e^{\frac{iv}{\beta}}e^{-\frac{y}{\beta}}e^{-\frac{i\pi}{\beta}}}.$$

Then it's easy to see that

$$F(v,y)=\frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]}.$$

Thus,

$$\int_{L_1+L_2} e^{z[\cos a]}\frac{1}{1-e^{-\frac{i(a-v)}{\beta}}}da = E(z,v).$$

Thus, after expanding we have that

$$P(z,v)=2\pi\beta\Sigma_{k,\ -\pi<v+2k\beta\pi<\pi}e^{z\cos(v+2k\beta\pi)}+\pi\beta\Sigma_{k,v+2k\beta\pi=\pm\pi}e^{-z}+E(z,v).$$

The proof is complete. □

Next we state the representation formulas for the second derivatives of the convolution of the heat kernel with a function, which are solutions to the parabolic equations with f as right hand side .

Lemma 4.5. *Suppose f is supported in A_{10}, then*

$$\begin{aligned}
&\frac{\partial}{\partial t}(H*f)(x)\\
= \ &f(x,t)+\int_0^t\int_{A_{10}}\frac{\partial}{\partial t}[H(x,y,t-\tau)][f(y,\tau)-f(x,\tau)]dyd\tau\\
&-\int_0^t f(x,\tau)d\tau\int_{\partial A_{10}} < \nabla_y H(x,y,t-\tau), n(y) > ds_y.
\end{aligned}$$

Furthermore, for any $1\le i\le m$ the following formulas are true.

$$\begin{aligned}
\frac{\partial^2}{\partial r\partial s_i}(H*f) \ = \ &\int_0^t\int_{A_{10}}\frac{\partial^2}{\partial r\partial s_i}[H(x,y,t-\tau)][f(y,\tau)-f(x,\tau)]dyd\tau\\
&-\int_0^t f(x,\tau)d\tau\int_{\partial A_{10}}\frac{\partial}{\partial r}H(x,y,t-\tau)n_i(y)ds_y;
\end{aligned}$$

$$\begin{aligned}\frac{1}{r}\frac{\partial^2}{\partial\theta\partial s_i}(H*f) &= \int_0^t\int_{A_{10}}\frac{1}{r}\frac{\partial^2}{\partial\theta\partial s_i}[H(x,y,t-\tau)][f(y,\tau)-f(x,\tau)]dyd\tau \\ &\quad -\int_0^t f(x,\tau)d\tau\int_{\partial A_{10}}\frac{1}{r}\frac{\partial}{\partial\theta}H(x,y,t-\tau)n_i(y)ds_y.\end{aligned}$$

n stands for the inward normal of ∂A_{10}.

Proof. of Lemma 4.5: We only prove the representation formula for $\frac{\partial^2}{\partial r\partial s_i}(H*f)$, the formula for $\frac{\partial}{\partial t}(H*f)$ and $\frac{1}{r}\frac{\partial^2}{\partial r\partial s_i}(H*f)$ are the same. It suffices to show that

$$\lim_{h\to 0}\frac{\partial^2}{\partial r\partial s_i}\int_0^t\int_{A_{10}}\frac{\partial^2}{\partial r\partial s_i}[H(x,y,t-\tau)\eta_h(y)]f(y,\tau)dyd\tau$$

exists and equals the right hand side, provided η_h is properly chosen.

The crucial point is that, we can choose a good η_h away from the singularity. Suppose x is not on the singular hypersurface $\{0\}\times\mathbb{R}^m$, there is a ball $B_h(x)$ of radius h which does not only miss $\{0\}\times\mathbb{R}^m$, but also has argument range less than $\beta\pi$. Then by the choice of $B_h(x)$, $d(x,y)$ is smooth in y in $B_h(x)$. We choose a smooth function ψ such that $\psi(u)=0$ when $u\leq\frac{1}{2}$; $\psi(u)=1$ when $u\geq 1$. Then by letting $\eta_h(y)=\psi(\frac{d(x,y)}{h})$, the proof is complete, as in [20]. ☐

The next lemma is obvious.

Proposition 4.6. *Suppose u is compactly supported in A_{10}. Suppose $u\in L^2[0,T;H_0^1(A_{10})]$ and $\frac{\partial u}{\partial t}\in L^2[0,T;H^{-1}(A_{10})]$. Suppose u solves $(\Delta-\frac{\partial}{\partial t})u=f$ in the weak sense over $A_{10}\times[0,T]$ and u has zero initial value. Suppose $f\in C^{\alpha,\beta}(A_{10}\times[0,T])$. Then*

$$u=-(H*f)(x,t)=-\int_0^t\int_{A_{10}}H(x,y,t-\tau)f(y,\tau)dyd\tau.$$

5 Hölder estimate of the singular integrals and proof of the main Schauder estimates in Theorem 1.8.

This section is devoted to the proof of Theorem 1.8, by establishing proposition 5.3, 5.4, 5.5, 5.6, 5.1, Theorem 5.7. All of these are achieved from the properties of the heat kernel in Theorem 1.11 (which will be proved in the later part of this article). However, for the sake of being precise we won't quote directly Theorem 1.11, but will appeal to the lemmas which Theorem 1.11 is consisted of. Notice that the estimate in this section are more or less standard, since we are equipped with Theorem 1.11.

Proposition 5.1. *Suppose u is compactly supported in A_1. Suppose $u \in L^2[0,T;H_0^1(A_1)]$ and $\frac{\partial u}{\partial t} \in L^2[0,T;H^{-1}(A_1)]$. Suppose u solves $(\Delta - \frac{\partial}{\partial t})u = f$ in the weak sense over $A_1 \times [0,T]$ and u has zero initial value. Suppose $f \in C^{\alpha,\frac{\alpha}{2},\beta}(A_1 \times [0,T])$. Then there exists a constant C as in Definition 2.6 such that the following estimates hold.*

$$[\sqrt{-1}\partial\bar{\partial}u]_{\alpha,\frac{\alpha}{2},\beta,A_1\times[0,T]} \leq C[f]_{\alpha,\frac{\alpha}{2},\beta,A_1\times[0,T]}$$

and

$$[\sqrt{-1}\frac{\partial}{\partial t}u]_{\alpha,\frac{\alpha}{2},\beta,A_1\times[0,T]} \leq C[f]_{\alpha,\frac{\alpha}{2},\beta,A_1\times[0,T]}.$$

Proof. of proposition 5.1:, By rescaling, this is a direct corollary of proposition 5.3, 5.4, 5.5, 5.6, and the representation formula in proposition 4.6. Just notice two points. First, we can take u to be supported in A_1 so we don't have to shrink domain. Second, by the same ideas in [17], all the terms in $\sqrt{-1}\partial\bar{\partial}u$ except the term $\Delta_\beta u$ are estimated by proposition 5.3, 5.4, 5.5, 5.6. Then to estimate $\Delta_\beta u$, it suffices to use the equation $(\Delta - \frac{\partial}{\partial t})u = f$ itself (as $\Delta = \Delta_\beta + \Sigma_{i=1}^m \frac{\partial^2}{\partial s_i^2}$). ☐

Corollary 5.2. *Suppose u is compactly supported in B_1. Suppose $u \in L^2[0,T;H_0^1(B_1)]$ and $\frac{\partial u}{\partial t} \in L^2[0,T;H^{-1}(B_1)]$. Suppose u solves $(\Delta - \frac{\partial}{\partial t})u = f$ in the weak sense over $B_1 \times [0,T]$ and u has zero initial value. Suppose $f \in C^{\alpha,\frac{\alpha}{2},\beta}(B_1 \times [0,T])$. Then there exists a constant C (as in Definition 2.6) such that the following estimates hold.*

$$[\sqrt{-1}\partial\bar{\partial}u]_{\alpha,\frac{\alpha}{2},\beta,B_1\times[0,T]} \leq C[f]_{\alpha,\frac{\alpha}{2},\beta,B_1\times[0,T]}$$

and

$$[\sqrt{-1}\frac{\partial}{\partial t}u]_{\alpha,\frac{\alpha}{2},\beta,B_1\times[0,T]} \leq C[f]_{\alpha,\frac{\alpha}{2},\beta,B_1\times[0,T]}.$$

Proof. of Corollary 5.2: It's actually straight forward to deduce this proposition using proposition 5.1. To fully illustrate the equivalence relation between B_1 and A_1, we shall include the detail here. We divide the situation into two cases in the following.

Case 1: When $dist\{B_1(p),\ \{0\}\times\mathbb{R}^m\} \geq 1000$, it's obvious since the equation is regular away from the singularity.

Case 2: When $dist\{B_1(p),\ \{0\}\times\mathbb{R}^m\}\leq 1000$, by translation invariance we can assume $\widehat{p}=0$. Then $B_1\subset A_{1100}$ and we have

$$\begin{aligned}
& [\sqrt{-1}\partial\bar{\partial}u]_{\alpha,\frac{\alpha}{2},\beta,B_1(p)\times[0,T]}\\
\leq\ & [\sqrt{-1}\partial\bar{\partial}u]_{\alpha,\frac{\alpha}{2},\beta,A_{1100}\times[0,T]}\\
\leq\ & [f]_{\alpha,\frac{\alpha}{2},\beta,A_{20000}\times[0,T]}\\
\leq\ & [f]_{\alpha,\frac{\alpha}{2},\beta,B_{30000}(p)\times[0,T]}.
\end{aligned}$$

Then since both u and f are supported in $B_1(p)$, the proof is completed. □

The next result is on estimates of spatial derivatives of the singular integral.

Proposition 5.3. *There exists a C (as in Definition 2.6) independent of T such that for any $\mathfrak{D}\in\mathfrak{T}$, the following estimate holds.*

$$[\mathfrak{D}(H*f)]_{\alpha,\beta,A_1\times[0,T]}\leq C[f]_{\alpha,\beta,A_{10}\times[0,T]}.$$

Proof. of proposition 5.3: The proof is similar to Donaldson's estimates in [17]. We only prove the statement for $\frac{\partial^2}{\partial r\partial s_i}$ with some fixed i. The statements for $\frac{\partial^2}{\partial s_i\partial s_j},\ i,j\leq m$ and $\frac{1}{r}\frac{\partial^2}{\partial s_i\partial\theta},\ i\leq m$ are proved in exactly the same way.

By rescaling it suffices to assume $|f|_{\alpha,\beta}=1$. Denote $|x_1-x_2|=\delta$.
Case1: $x_1=0$ (or $x_2=0$) , we have

$$\begin{aligned}
& \frac{\partial^2}{\partial r\partial s_i}(H*f)(0,t)-\frac{\partial^2}{\partial r\partial s_i}(H*f)(x_2,t)\\
=\ & I_1+I_2+I_3+I_4+I_5+I_6,
\end{aligned}$$

where

$$\begin{aligned}
I_1 &= \int_0^t \int_{\{|y|\le 64\delta\}\cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(0,y,t-\tau)][f(y,\tau)-f(0,\tau)]dyd\tau;\\
I_2 &= -\int_0^t \int_{\{|y|\le 64\delta\}\cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_2,y,t-\tau)][f(y,\tau)-f(x_2,\tau)]dyd\tau;\\
I_3 &= \int_0^t \int_{\{|y|\ge 64\delta\}\cap A_{10}} [\frac{\partial^2}{\partial r \partial s_i}H(0,y,t-\tau)-\frac{\partial^2}{\partial r \partial s_i}H(x_2,y,t-\tau)]\}\\
&\quad \times[f(y,\tau)-f(0,\tau)]dyd\tau;\\
I_4 &= \int_0^t [f(x_2,\tau)-f(0,\tau)]d\tau \int_{\{|y|\ge 64\delta\}\cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_2,y,t-\tau)]dy;\\
I_5 &= -\int_0^t [f(0,\tau)-f(x_2,\tau)]d\tau \int_{\partial A_{10}} [\frac{\partial}{\partial r}H(0,y,t-\tau)]n_i(y)ds_y;\\
I_6 &= -\int_0^t f(x_2,\tau)d\tau \int_{\partial A_{10}} [\frac{\partial}{\partial r}H(0,y,t-\tau)-\frac{\partial}{\partial r}H(x_2,y,t-\tau)]n_i(y)ds_y.
\end{aligned}$$

On I_1, by Item 1 in Lemma 7.1, we have $\int_0^{+\infty}|\frac{\partial^2}{\partial r\partial s_i}H(0,y,t)|dt \le C|y|^{-(m+2)}$. Then

$$|I_1| \le C\int_{\{|y|\le 64\delta\}\cap A_{10}} |y|^{-(m+2)+\alpha}dy \le C\delta^\alpha.$$

On I_2, by item 2 in Lemma 7.1 and rescaling, we have

$$\int_0^{+\infty}|\frac{\partial^2}{\partial r\partial s_i}H(x_2,y,t)|dt \le C|y-x_2|^{-(m+2)}.$$

Thus

$$|I_2| \le C\int_{\{|y|\le 64\delta\}\cap A_{10}} |y-x_2|^{-(m+2)+\alpha}dy \le C\delta^\alpha.$$

On I_3, by Lemma 7.3 and rescaling, we have

$$\int_0^{+\infty}|\frac{\partial^2}{\partial r\partial s_i}H(x_2,y,t)-\frac{\partial^2}{\partial r\partial s_i}H(0,y,t)|dt \le C\delta|y|^{-(m+3)}.$$

Then

$$|I_3| \le C\delta\int_{\{|y|\ge 64\delta\}\cap A_{10}} |y|^{-(m+2)-1+\alpha}dy \le C\delta^\alpha.$$

On I_4, by Lemma 7.2 we have $\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau \le C\delta^{-(m+1)}$ when $|x_2| = \delta$ and $|y| = 64\delta$. With the conclusion of Lemma 7.2 in mind, we compute that

$$
\begin{aligned}
& |I_4| \\
\leq\ & \int_0^t |f(x_2,\tau)-f(0,\tau)|d\tau \int_{|y|=64\delta} |\frac{\partial}{\partial r}[H(x_2,y,t-\tau)|ds_y \\
& + \int_0^t |f(x_2,\tau)-f(0,\tau)|d\tau \int_{\partial A_{10}} |\frac{\partial}{\partial r}[H(x_2,y,t-\tau)|ds_y \\
\leq\ & C\delta^\alpha[\int_{|y|=64\delta}\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau ds_y + \int_{\partial A_{10}}\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau ds_y] \\
\leq\ & C\delta^\alpha.
\end{aligned}
$$

On I_5, we have

$$
\begin{aligned}
& |\int_0^t [f(0,\tau)-f(x_2,\tau)]d\tau \int_{\partial A_{10}} < \frac{\partial}{\partial r}H(0,y,t-\tau), n_i(y) > ds_y| \\
\leq\ & \delta^\alpha \int_0^t d\tau \int_{\partial A_{10}} |\frac{\partial}{\partial r}H(0,y,t-\tau), n_i(y)|ds_y.
\end{aligned}
$$

Still by Lemma 7.2, we have over ∂A_{10} that

$$
\int_0^t |\frac{\partial}{\partial r}H(0,y,t-\tau), n_i(y)|d\tau \leq C,
$$

then $|I_5| \leq C\delta^\alpha$.

The estimate of I_6 is the reason we use the polydisk A_{10} but not balls. Recall from Definition that

$$
\partial A_{10} = [\partial D_{10} \times \widehat{B}_{10}] \cup [D_{10} \times \partial \widehat{B}_{10}].
$$

Over $\partial D_{10} \times \widehat{B}_{10}$, we have $n_i(y) = 0$. Therefore the second inequality in Lemma 8.2 directly implies that

$$
\begin{aligned}
& \int_0^t d\tau \int_{\partial A_{10}} |\frac{\partial}{\partial r}H(0,y,t-\tau) - \frac{\partial}{\partial r}H(x_2,y,t-\tau), n_i(y)|ds_y \\
\leq\ & \int_{D_{10}\times\partial\widehat{B}_{10}} \int_0^{+\infty} |\frac{\partial}{\partial r}H(0,y,t-\tau) - \frac{\partial}{\partial r}H(x_2,y,t-\tau)|d\tau ds_y \\
\leq\ & C\delta^\rho,
\end{aligned}
$$

which means $|I_6| \leq C\delta^\rho$. Thus the proof is completed in this case.

Case 2: We assume $\min\{P(x_1), P(x_2)\} = l \geq 3\delta$, where $P(x)$ is norm of the projection of x to the $\mathbb{R}^2$ component. This case is easier to handle since both points are far away from the singularity. Without loss of generality we can assume $\widehat{x_1} = 0$ and then $|x_1| = l\delta$. Then we split the difference as

$$\begin{aligned}
&\frac{\partial^2}{\partial r \partial s_i}(H * f)(x_1, t) - \frac{\partial^2}{\partial r \partial s_i}(H * f)(x_2, t) \\
= \ & IV_1 + IV_2 + IV_3 + IV_4 + IV_5 + IV_6 + IV_7 + IV_8,
\end{aligned}$$

where

$$\begin{aligned}
IV_1 &= \int_0^t \int_{\{|y| \leq 2l\delta, |y - x_1| \leq 64\delta\} \cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_1, y, t - \tau)][f(y, \tau) - f(x_1, \tau)] dy d\tau; \\
IV_2 &= -\int_0^t \int_{\{|y| \leq 2l\delta, |y - x_1| \leq 64\delta\} \cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_2, y, t - \tau)][f(y, \tau) - f(x_2, \tau)] dy d\tau; \\
IV_3 &= \int_0^t \int_{\{|y| \leq 2l\delta, |y - x_1| \geq 64\delta\} \cap A_{10}} [\frac{\partial^2}{\partial r \partial s_i} H(x_1, y, t - \tau) - \frac{\partial^2}{\partial r \partial s_i} H(x_2, y, t - \tau)]\} \\
&\quad \times [f(y, \tau) - f(x_1, \tau)] dy d\tau; \\
IV_4 &= \int_0^t [f(x_2, \tau) - f(x_1, \tau)] d\tau \int_{\{|y| \leq 2l\delta, |y - x_1| \geq 64\delta\} \cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_2, y, t - \tau)] dy; \\
IV_5 &= \int_0^t \int_{\{|y| \geq 2l\delta\} \cap A_{10}} [\frac{\partial^2}{\partial r \partial s_i} H(x_1, y, t - \tau) - \frac{\partial^2}{\partial r \partial s_i} H(x_2, y, t - \tau)]\} \\
&\quad \times [f(y, \tau) - f(x_1, \tau)] dy d\tau; \\
IV_6 &= \int_0^t [f(x_2, \tau) - f(x_1, \tau)] d\tau \int_{\{|y| \geq 2l\delta\} \cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_2, y, t - \tau)] dy; \\
IV_7 &= \int_0^t [f(x_1, \tau) - f(x_2, \tau)] d\tau \int_{\partial A_{10}} < \frac{\partial}{\partial r} H(x_1, y, t - \tau), n_i(y) > ds_y; \\
IV_8 &= \int_0^t f(x_2, \tau) d\tau \int_{\partial A_{10}} < \frac{\partial}{\partial r} H(x_1, y, t - \tau) - \frac{\partial}{\partial r} H(x_2, y, t - \tau), n_i(y) > ds_y.
\end{aligned}$$

Mostly we will proceed as in Case 1, except on some special items.

On IV_1, from the same analysis as in I_2 in Case 1, we have

$$\begin{aligned}
|IV_1| &\leq \int_0^t \int_{\{|y - x_1| \leq 64\delta\} \cap A_{10}} \frac{\partial^2}{\partial r \partial s_i}[H(x_1, y, t - \tau)][f(y, \tau) - f(x_1, \tau)] dy d\tau \\
\leq \ & C \int_{\{|y - x_1| \leq 64\delta\} \cap A_{10}} |y - x_1|^{-(m+2)+\alpha} dy \leq C\delta^\alpha.
\end{aligned}$$

Second we compute

$$|IV_2| \leq \int_0^t \int_{\{|y-x_2|\leq 65\delta\}\cap A_{10}} \frac{\partial^2}{\partial r\partial s_i}[H(x_2,y,t-\tau)][f(y,\tau)-f(x_2,\tau)]dyd\tau$$

$$\leq \;\; C\int_{\{|y-x_2|\leq 65\delta\}\cap A_{10}} |y-x_2|^{-(m+2)+\alpha}dy \leq C\delta^\alpha.$$

On IV_3, notice that when $|y|\leq 2l\delta$ and $|y-x|\geq 63\delta$, by Lemma 7.5, we have

$$\int_0^\infty |\nabla_x \frac{\partial^2}{\partial r\partial s_i}H(x,y,\tau)|d\tau \leq C|y-x|^{-(m+3)}$$

for any x in the line segment $\overline{x_1x_2}$. Thus

$$\begin{aligned}&|IV_3|\\ \leq\;\; & C\delta\int_{\{|y-x_1|\geq 64\delta\}\cap A_{10}} |y-x_1|^{-(m+3)+\alpha}dy \leq C\delta^\alpha.\end{aligned}$$

On IV_4, similarly we have

$$\begin{aligned}&|IV_4|\\ \leq\;\; & C\delta^\alpha\int_0^t\int_{|y|=2l\delta}|\frac{\partial}{\partial r}[H(x_2,y,t-\tau)]|d\tau ds(y)\\ &+C\delta^\alpha\int_0^t\int_{|y-x_1|=64\delta}|\frac{\partial}{\partial r}[H(x_2,y,t-\tau)]|d\tau ds(y)\\ &+C\delta^\alpha\int_0^t\int_{\partial A_{10}}|\frac{\partial}{\partial r}[H(x_2,y,t-\tau)]|d\tau ds(y).\end{aligned}$$

By Lemma 7.2 we have

$$\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau \leq C|y-x_2|^{-(m+1)}$$

when $|y-x_1|=64\delta$. Therefore when $|y|=2l\delta$, since $l\geq 2$, we have

$$\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau \leq C|y-x_2|^{-(m+1)} \leq C\delta^{-(m+1)},$$

thus

$$\int_{|y|=2l\delta}\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau ds(y) \leq C.$$

When $|y-x_1|=64\delta$, we also have

$$\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau \leq C|y-x_2|^{-(m+1)} \leq C\delta^{-(m+1)},$$

thus

$$\int_{|y-x_1|=64\delta}\int_0^\infty |\frac{\partial}{\partial r}[H(x_2,y,\tau)]|d\tau ds(y) \le C.$$

According to Lemma 7.2 and the assumption that $|x_2| \le 1$, we have

$$\int_0^t \int_{\partial A_{10}} |\frac{\partial}{\partial r}[H(x_2,y,t-\tau)]|d\tau ds(y) \le C.$$

Then $|IV_4| \le C\delta^\alpha$.

IV_6 can be estimated easily in the way IV_4 was estimated. IV_5 can be estimated exactly as I_3 in Case 1. The estimates of IV_7 and IV_8 are exactly the same as the estimates of I_5 and I_6. Thus case 2 is all set.

Finally, given any $x_1,\ x_2$ such that $|x_1 - x_2| = \delta$, we always have

$$|\frac{\partial^2}{\partial r\partial s_i}(H*f)(x_1,t) - \frac{\partial^2}{\partial r\partial s_i}(H*f)(x_2,t)| \le C\delta^\alpha$$

The reason is that by translation invariance we can assume $\widehat{x_1} = 0$ and $P(x_1) \le P(x_2)$. If $P(x_1) \le 3\delta$, we have:

$$\begin{aligned} & |\frac{\partial^2}{\partial r\partial s_i}(H*f)(x_1,t) - \frac{\partial^2}{\partial r\partial s_i}(H*f)(x_2,t)| \\ \le\ & |\frac{\partial^2}{\partial r\partial s_i}(H*f)(x_1,t) - \frac{\partial^2}{\partial r\partial s_i}(H*f)(0,t)| \\ & +|\frac{\partial^2}{\partial r\partial s_i}(H*f)(x_2,t) - \frac{\partial^2}{\partial r\partial s_i}(H*f)(0,t)| \\ \le\ & C\delta^\alpha \end{aligned}$$

from Case 1. If $P(x_1) \ge 3\delta$ we are in Case 2. Thus the proof is completed. □

Proposition 5.4. *There exists a C (as in Definition 2.6) independent of T such that*

$$[\frac{\partial}{\partial t}(H*f)]_{\alpha,\beta,A_1\times[0,T]} \le C[f]_{\alpha,\beta,A_{10}\times[0,T]}.$$

Proof. of proposition 5.4: The framework of this lemma is simpler than that of Lemma 5.3. We will repeatedly use the assumptions $|x_1|, |x_2| \le 1$. By rescaling it suffices to assume $|f|_{\alpha,\beta} = 1$. Denote $|x_1 - x_2| = \delta$. We compute

$$\begin{aligned} & \frac{\partial}{\partial t}(H*f)(x_1,t) - \frac{\partial}{\partial t}(H*f)(x_2,t) \\ =\ & f(x_1,t) - f(x_2,t) + Z_1 + Z_2 + Z_3 + Z_4 + Z_5 + Z_6. \end{aligned}$$

$$
\begin{aligned}
Z_1 &= \int_0^t \int_{\{|y-x_1|\le 64\delta\}\cap A_{10}} \frac{\partial}{\partial t}[H(x_1,y,t-\tau)][f(y,\tau)-f(x_1,\tau)]dyd\tau;\\
Z_2 &= -\int_0^t \int_{\{|y-x_1|\le 64\delta\}\cap A_{10}} \frac{\partial}{\partial t}[H(x_2,y,t-\tau)][f(y,\tau)-f(x_2,\tau)]dyd\tau;\\
Z_3 &= \int_0^t \int_{\{|y-x_1|\ge 64\delta\}\cap A_{10}} [\frac{\partial}{\partial t}H(x_1,y,t-\tau)-\frac{\partial}{\partial t}H(x_2,y,t-\tau)]\}\\
&\quad [f(y,\tau)-f(x_1,\tau)]dyd\tau;\\
Z_4 &= \int_0^t [f(x_2,\tau)-f(x_1,\tau)]d\tau \int_{\{|y-x_1|\ge 64\delta\}\cap A_{10}} \frac{\partial}{\partial t}[H(x_2,y,t-\tau)]dy;\\
Z_5 &= -\int_0^t [f(x_1,\tau)-f(x_2,\tau)]d\tau \int_{\partial A_{10}} <\nabla_y H(x_1,y,t-\tau),n(y)> ds_y;\\
Z_6 &= -\int_0^t f(x_2,\tau)d\tau \int_{\partial A_{10}} <\nabla_y H(x_1,y,t-\tau)-\nabla_y H(x_2,y,t-\tau),n(y)> ds_y.
\end{aligned}
$$

On Z_1, from Lemma 7.1, we have $\int_0^{+\infty}|\frac{\partial}{\partial t}H(x_1,y,t)|dt \le C|y-x_1|^{-(m+2)}$. Then

$$|Z_1| \le C\int_{\{|y-x_1|\le 64\delta\}\cap A_{10}} |y|^{-(m+2)+\alpha}dy \le C\delta^\alpha.$$

On Z_2, again from Lemma 7.1 and rescaling, we have

$$\int_0^{+\infty}|\frac{\partial}{\partial t}H(x_2,y,t)|dt \le C|y-x_2|^{-(m+2)}.$$

Thus

$$|Z_2| \le C\int_{\{|y-x_1|\le 64\delta\}\cap A_{10}} |y-x_2|^{-(m+2)+\alpha}dy \le C\delta^\alpha.$$

On Z_3, from Lemma 7.3 and rescaling, we have

$$\int_0^{+\infty}|\frac{\partial}{\partial t}H(x_2,y,t)-\frac{\partial}{\partial t}H(x_1,y,t)|dt \le C\delta|y-x_1|^{-(m+3)}.$$

Then

$$|Z_3| \le C\delta\int_{\{|y-x_1|\ge 64\delta\}\cap A_{10}} |y-x_1|^{-(m+2)-1+\alpha}dy \le C\delta^\alpha.$$

On Z_4, since we have $\int_0^\infty |\nabla_y H(x_2,y,\tau)|d\tau \le C\delta^{-(m+1)}$ for $|x_2|=\delta$ and

$|y-x_1|=64\delta$, then

$$\begin{aligned}
& |Z_4| \\
= & \ |\int_0^t [f(x_2,\tau)-f(x_1,\tau)]d\tau \int_{\{|y-x_1|\geq 64\delta\}\cap A_{10}} \Delta[H(x_2,y,t-\tau)]dy| \\
\leq & \ |\int_0^t [f(x_2,\tau)-f(x_1,\tau)]d\tau \int_{\partial[\{|y-x_1|=64\delta\}\cap A_{10}]} \\
& < \nabla_y[H(x_2,y,t-\tau)], n(y) > ds_y| \\
\leq & \ C\delta^\alpha [\int_{|y-x_1|=64\delta} \int_0^\infty |\nabla_y[H(x_2,y,\tau)]| d\tau ds_y \\
& + \int_{\partial A_{10}} \int_0^\infty |\nabla_y[H(x_2,y,\tau)]| d\tau ds_y] \\
\leq & \ C\delta^\alpha .
\end{aligned}$$

On Z_5, we have

$$\begin{aligned}
& |\int_0^t [f(x_1,\tau)-f(x_2,\tau)]d\tau \int_{\partial A_{10}} < \nabla_y H(x_1,y,t-\tau), n(y) > ds_y| \\
\leq & \ \delta^\alpha \int_0^t d\tau \int_{\partial A_{10}} |\nabla_y H(x_1,y,t-\tau)| ds_y .
\end{aligned}$$

Still by Lemma 7.2, we have over ∂A_{10} that

$$\int_0^t |\nabla_y H(x_1,y,t-\tau))| d\tau \leq C,$$

then $|Z_5| \leq C\delta^\alpha$.

The estimate of Z_6 is the reason we use the polydisk A_{10} but not balls. Again recall from the definitions that

$$\partial A_{10} = [\partial D_{10} \times \widehat{B}_{10}] \cup [D_{10} \times \partial \widehat{B}_{10}].$$

Over $\partial D_{10} \times \widehat{B}_{10}$, we have $n(y) = \frac{\partial}{\partial \acute{r}}$; Over $D_{10} \times \partial \widehat{B}_{10}$, we have $n(y) = \frac{\partial}{\partial R}$. The easy but crucial observation is that the normal vectors are all orthogonal to $\frac{\partial}{\partial \acute{\theta}}$. Therefore Lemma 7.6 directly implies that

$$
\begin{aligned}
& \int_0^t d\tau \int_{\partial A_{10}} |\nabla_y H(x_1, y, t-\tau) - \nabla_y H(x_2, y, t-\tau), n(y)| ds_y \\
\leq \; & \delta \int_0^t d\tau \int_{\partial D_{10} \times \widehat{B}_{10}} |\nabla_{\bar{x}} \frac{\partial}{\partial \acute{r}} H[\bar{x}(y, t-\tau), y, t-\tau]| ds_y \\
& + \delta \int_0^t d\tau \int_{(D_{10} \setminus \{0\}) \times \partial \widehat{B}_{10}} |\nabla_{\bar{x}} \frac{\partial}{\partial \bar{R}} H[\bar{x}(y, t-\tau), y, t-\tau]| ds_y \\
\leq \; & C\delta,
\end{aligned}
$$

which means $|Z_6| \leq C\delta$. Thus the proof is done in this case.

$\square$

In the following we shall study the timewise Hölder estimate of the spatial second derivatives and time derivative of the solution. The main conclusions in section 9 play important roles here.

Proposition 5.5. *There exists a C (as in Definition 2.6) with the following property. For all second order differential operators $\mathfrak{D} \in \mathfrak{T}$, points $x \in A_1$ and t such that $0 \leq t < t + \delta \leq T$, the following potential estimate holds.*

$$
|\mathfrak{D}(H * f)(x, t+\delta) - \mathfrak{D}(H * f)(x, t)| \leq C\delta^{\frac{\alpha}{2}} [f]_{\alpha, \beta, A_{10} \times [0, T]}.
$$

Proof. of Proposition 5.5: It suffices to assume $\delta \leq \frac{1}{10000}$ in the Hölder estimates. The case when $\delta \geq \frac{1}{10000}$ reduces to lower order estimates in section 11 which are a lot easier than the estimates we try to prove here. The necessary lower order estimates are stated in section 11. In the estimate of Q_4 below, we applied Lemma 9.1 which requires the δ to be small.

By rescaling invariance of the heat kernel it suffices to assume $|f|_{\alpha,\beta} = 1$. It suffices to only consider $\mathfrak{D} = \frac{\partial^2}{\partial r \partial s_i}$, since the other derivatives are really similar. We compute

$$
\begin{aligned}
& (\frac{\partial^2}{\partial r \partial s_i} H * f)(x, t+\delta) - (\frac{\partial^2}{\partial r \partial s_i} H * f)(x, t) \\
= \; & Q_1 + Q_2 + Q_3 + Q_4 + Q_5,
\end{aligned}
$$

where

$$
\begin{aligned}
Q_1 &= \int_{t-\delta}^{t+\delta} d\tau \int_{A_{10}} \frac{\partial^2}{\partial r \partial s_i} H(x,y,t+\delta-\tau)[f(y,\tau)-f(x,\tau)]dy \\
Q_2 &= -\int_{t-\delta}^{t} d\tau \int_{A_{10}} \frac{\partial^2}{\partial r \partial s_i} H(x,y,t-\tau)][f(y,\tau)-f(x,\tau)]dy \\
Q_3 &= \int_{0}^{t-\delta} d\tau \int_{A_{10}} [\frac{\partial^2}{\partial r \partial s_i} H(x,y,t+\delta-\tau) - \frac{\partial^2}{\partial r \partial s_i} H(x,y,t-\tau)] \\
\times & \quad [f(y,\tau)-f(x,\tau)]dy \\
Q_4 &= -\int_{t}^{t+\delta} f(x,\tau) d\tau \int_{\partial A_{10}} \frac{\partial}{\partial r} H(x,y,t+\delta-\tau) n_i(y) ds_y \\
Q_5 &= -\int_{0}^{t} f(x,\tau) d\tau \int_{\partial A_{10}} [\frac{\partial}{\partial r} H(x,y,t+\delta-\tau) - \frac{\partial}{\partial r} H(x,y,t-\tau)] n_i(y) ds_y
\end{aligned}
$$

Next, using the estimates in Proposition 9.2, we estimate that

$$
\int_{\mathbb{R}^2 \times \mathbb{R}^m} |\frac{\partial^2}{\partial r \partial s_i} H(x,y,t)||y-x|^\alpha dy \le C t^{\frac{\alpha}{2}-1},
$$

then we estimate

$$
\begin{aligned}
& |Q_1| \\
\le & \int_0^{2\delta} dt \int_{\mathbb{R}^2 \times \mathbb{R}^m} |\frac{\partial^2}{\partial r \partial s_i} H(x,y,t)||y-x|^\alpha dy \\
\le & C \int_0^{2\delta} t^{\frac{\alpha}{2}-1} dt \\
\le & C \delta^{\frac{\alpha}{2}}.
\end{aligned}
$$

Similarly we have $|Q_2| \le C\delta^{\frac{\alpha}{2}}$.

For Q_3, we compute

$$
\begin{aligned}
& |Q_3| \\
\le\ & \int_{-\infty}^{t-\delta} d\tau \int_{A_{10}} |\frac{\partial^2}{\partial r \partial s_i} H(x,y,t+\delta-\tau) - \frac{\partial^2}{\partial r \partial s_i} H(x,y,t-\tau)|\ |y-x|^{\alpha} dy \\
=\ & \int_{\delta}^{\infty} t^{\frac{\alpha}{2}-1} dt \int_{\frac{A_{10}}{t}} |\frac{\partial^2}{\partial r \partial s_i} H(\frac{x}{\sqrt{t}}, u, 1+\frac{\delta}{t}) - \frac{\partial^2}{\partial r \partial s_i} H(\frac{x}{\sqrt{t}}, u, 1)|\ |u-\frac{x}{\sqrt{t}}|^{\alpha} du \\
\le\ & C\delta \int_{\delta}^{\infty} t^{\frac{\alpha}{2}-2} dt \int_{R^2 \times R^m} \sup_{1 \le s \le 2} |\frac{\partial}{\partial s} \frac{\partial^2}{\partial r \partial s_i} H(\frac{x}{\sqrt{t}}, u, s)||u-\frac{x}{\sqrt{t}}|^{\alpha} du \\
\le\ & C\delta \int_{\delta}^{\infty} t^{\frac{\alpha}{2}-2} dt \\
\le\ & C\delta^{\frac{\alpha}{2}},
\end{aligned}
$$

where we applied the following estimate in Proposition 9.2:

$$
\int_{\mathbb{R}^2 \times \mathbb{R}^m} \sup_{1 \le s \le 2} |\frac{\partial}{\partial s} \frac{\partial^2}{\partial r \partial s_i} H(\frac{x}{\sqrt{t}}, u, s)||u-\frac{x}{\sqrt{t}}|^{\alpha} du \le C.
$$

On Q_4, we have to involve Lemma 9.1. We compute

$$
\begin{aligned}
& |Q_4| \\
=\ & \int_0^{\delta} dt \int_{\partial A_{10}} |\frac{\partial}{\partial r} H(x,y,t)| ds_y \\
=\ & \int_0^{\delta} t^{-1} dt \int_{\frac{\partial A_{10}}{t}} |\frac{\partial}{\partial r} H(\frac{x}{t}, u, 1)| ds_u \le \int_0^{\delta} t^{-1} \cdot t^{-\frac{m+1}{2}} e^{-\frac{1}{t}} dt \\
\le\ & C\delta.
\end{aligned}
$$

On Q_5, from Lemma 7.3, we have

$$
\begin{aligned}
& |Q_5| \\
\le\ & \delta \int_0^t d\tau \int_{\partial A_{10}} |\frac{\partial^2}{\partial r \partial t} H(x,y,\bar{t}(x,y)-\tau)| ds_y \\
\le\ & C\delta,
\end{aligned}
$$

where $t \le \bar{t}(x,y) \le t+\delta$.

□

Proposition 5.6. *There exists a C (as in Definition 2.6) with the following properties. For all points $x \in A_1$ and t_1, t_2 in $[0,\ T]$, the following potential estimate holds.*

$$
|(\frac{\partial H}{\partial t} * f)(x,t_1) - (\frac{\partial H}{\partial t} * f)(x,t_2)| \le C[f]_{\alpha,\frac{\alpha}{2},\beta,A_{10}\times[0,T]}|t_1-t_2|^{\frac{\alpha}{2}}.
$$

Proof. of Proposition 5.6: The proof is by the same way as in Lemma 5.5. We apply the following decomposition after rescaling with respect time and applying proposition 9.2.

$$\begin{aligned}
& \frac{\partial H}{\partial t} * f(x,t+\delta) - \frac{\partial H}{\partial t} * f(x,t) \\
= \; & (\frac{\partial H}{\partial t} H * f)(x,t+\delta) - (\frac{\partial H}{\partial t} H * f)(x,t) \\
= \; & \int_{t-\delta}^{t+\delta} d\tau \int_{A_{10}} \frac{\partial H}{\partial t} H(x,y,t+\delta-\tau)[f(y,\tau)-f(x,\tau)]dy \\
& - \int_{t-\delta}^{t} d\tau \int_{A_{10}} \frac{\partial H}{\partial t} H(x,y,t-\tau)][f(y,\tau)-f(x,\tau)]dy \\
& + \int_{-\infty}^{t-\delta} d\tau \int_{A_{10}} [\frac{\partial H}{\partial t} H(x,y,t+\delta-\tau) - \frac{\partial H}{\partial t} H(x,y,t-\tau)] \\
& \times [f(y,\tau)-f(x,\tau)]dy \\
& - \int_{t}^{t+\delta} f(x,\tau)d\tau \int_{\partial A_{10}} < \nabla_y H(x,y,t+\delta-\tau), \ n(y) > ds_y \\
& - \int_{0}^{t} f(x,\tau)d\tau \int_{\partial A_{10}} < \nabla_y [H(x,y,t+\delta-\tau) - H(x,y,t-\tau)], n_i(y) > ds_y \\
& + f(x,t+\delta) - f(x,t).
\end{aligned}$$

□

The next theorem on interior estimates is important. As we mentioned earlier, the singular set is never considered as boundary.

Theorem 5.7. *There exists a $C(T)$ (as in Definition 2.6 and depending on T) with the following properties. For any open set $\Omega \in \mathbb{R}^{m+2}$, suppose $u \in C_0^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$. Then u satisfies the following estimates*

$$\begin{aligned}
& |u|^{(\star)}_{2+\alpha,1+\frac{\alpha}{2},\beta,\Omega\times[0,T]} \leq C(T)|(\Delta - \frac{\partial}{\partial t})u|^{(2)}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]}; \\
& |u|^{(\star)}_{2,\alpha,\beta,\Omega\times[0,T]} \leq C(T)|(\Delta - \frac{\partial}{\partial t})u|^{(2)}_{\alpha,\beta,\Omega\times[0,T]}; \\
& |u|^{[\star]}_{2+\alpha,1+\frac{\alpha}{2},\beta,\Omega\times[0,T]} \leq C(T)|(\Delta - \frac{\partial}{\partial t})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]}.
\end{aligned}$$

Proof. of Theorem 5.7: Using Proposition 5.1, multiplication by a cutoff function, and the lower order estimates in the appendix, it's routine to deduce the estimates in Theorem 5.7. We should apply Proposition 5.1 near the singularities and apply Proposition 5.2 away from the singularities. Nevertheless, we would like to mention that the interpolation tricks we implicitly employed here are not the same ones as in [20], although the tricks are extremely simple and natural. Notice that away from the singularity we should apply Corollary 5.2. □

To prove the part of Theorem 1.8 on existence of classical solutions, we need the following existence result over E_T. The power of the space $W_0^{2,2}(E_T)$ is displayed here. The following result is definitely true when $\beta = 1$ (when there is no singularity), and we believe this is more than well known to experts.

Proposition 5.8. *The parabolic operator* $\Delta_{\omega_D} - \frac{\partial}{\partial t}$ *is surjective map among the following function spaces*

$$C_0^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T] :\rightarrow\ C^{\alpha,\frac{\alpha}{2},\beta}[0,T].$$

Proof. of Proposition 5.8: Though the aprori estimates in Theorem 5.7 hold, it's still necessary to show the existence of a classical solution. Since Proposition 5.8 has its own independent interest and needs the estimates in $W^{2,2}(M \times [0,T])$ (which involve a bunch of other techniques), we refer the reader to the note [48]. □

Now we begin addressing the final step of the main Schauder estimates and existence results in Theorem 1.8, by using the results established earlier in this section.

Proof. of Theorem 1.8: First we show the aprori estimates hold. Namely, suppose $u \in C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$, we prove that the estimates in Theorem 1.8 hold. These go exactly the same as in [20]. Notice that away from the singularity we have no problem at all, then what we need to prove more than Theorem 5.7 is that we should show the Schauder estimate for the space-time dependent operator $\Delta_{a(t)} - \frac{\partial}{\partial t}$, not only for the constant coefficient operator $\Delta - \frac{\partial}{\partial t}$ in Theorem 5.7.

We include the crucial detail on how to deal with the space-time dependent operator. We would like to point out that though the intepolation inequalities which are hiding behind should be proved using some different method, these estimates here are absolutely regular. Let h be small enough with respect to g, by Theorem 5.7 and the discussion in section 4.2 of [17], there exists a D-diagonal constant coeffificient operator Δ_{a_0} such that

$$\begin{aligned}
&|i\partial\bar{\partial}u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}\\
\leq\ & C|(\Delta_{a_0}-\frac{\partial}{\partial t})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}+C|u|_{0,A_h\times[t,t+h^2]}\\
\leq\ & C|(\Delta_{a(t)}-\frac{\partial}{\partial t})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}+C|(\Delta_{a(t)}-\Delta_{a_0})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}\\
&+C|u|_{0,A_h\times[t,t+h^2]}.
\end{aligned}$$

By further shrinking h to be small enough with respect to $a(t)$, we have that

$$\begin{aligned}
&C|(\Delta_{a(t)}-\Delta_{a_0})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}\\
\leq\ & \frac{1}{4}[i\partial\bar{\partial}u]^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}+Ch^{\alpha}[i\partial\bar{\partial}u]^{[2]}_{0,A_h\times[t,t+h^2]}.
\end{aligned}$$

Then by making h even smaller, we arrive at

$$[i\partial\bar{\partial}u]^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}\leq C|(\Delta_{a(t)}-\frac{\partial}{\partial t})u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[t,t+h^2]}+C|u|_{0,A_h\times[t,t+h^2]}. \tag{21}$$

As the above, by making h smaller if necessary, we get

$$|i\partial\bar{\partial}u]^{(2)}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[0,h^2|}\leq C|(\Delta_{a(t)}-\frac{\partial}{\partial t})u|^{(2)}_{\alpha,\frac{\alpha}{2},\beta,A_h\times[0,h^2]}+C|u|_{0,A_h\times[0,h^2]}. \tag{22}$$

Combing (21), (22), Corollary 5.2, and the C^0 estimate in Lemma 11.4, we get

$$|i\partial\bar{\partial}u|_{\alpha,\frac{\alpha}{2},\beta,M\times[0,T]}\leq C_a|v|_{\alpha,\frac{\alpha}{2},\beta,M\times[0,T]}.$$

The rest of the story are all of lower order, which is easier and regular. See in [20], [27],[25].

Second, the existence of a classical solution follows from deforming the solution of the parabolic equation of the operator $\Delta_{\omega_D}-\frac{\partial}{\partial t}$ (ω_D as in (1)). The invertibleness of $\Delta_{\omega_D}-\frac{\partial}{\partial t}$ is guaranteed by Proposition 5.8. The existence of the deformation is directly implied by the aprori estimates established above and a generalization of Theorem 5.2 in [20] (on existence of continuous deformation of solutions).

The proof of Theorem 1.8 is now complete. □

6 Heat kernel and Bessel functions of the second kind.

In this section, we use Bessel functions of the second kind to give a shorter proof of the estimates on integral of Bessel functions used in [17]. The results here are slightly more general than those in [17]. First, a lemma on Bessel function of the second kind.

Lemma 6.1. *For any* $\epsilon > 0$*, nonnegative integer* N*, and exponents* $\mu_1, \mu_2 \geq 0$*, such that* $\frac{N}{\beta} - \mu_1 \geq 0$*, there is a constant* $C(\epsilon, N, \mu_1, \mu_2)$ *(as in Definition 2.6 and might tend to* ∞ *as* ϵ *goes to* 0*) such that the following estimate on the weighted summation of Bessel functions holds.*

$$\Sigma_{k=N}^{\infty} k^{\mu_2} I_{\frac{k}{\beta}-\mu_1}(z) \leq C(\epsilon, N, \mu_1, \mu_2) z^{\frac{N}{\beta}-\mu_1} e^{\frac{3z}{2}} e^{\epsilon z}.$$

Proof. of Lemma 6.1: The estimate directly follows from the property of Gamma functions. We only prove the case when $N = 1,\ \mu_2 = 2,\ \mu_1 = 0$ i.e $\Sigma_{k=1}^{\infty} k^2 I_{\frac{k}{\beta}}(z) \leq C(\epsilon) z^{\frac{1}{\beta}} e^{\frac{3z}{2}} e^{\epsilon z}$. The rest follows similarly.

Again from section (7.25) of [49] we have

$$I_{\frac{k}{\beta}}(z) \leq \frac{\sqrt{\pi} z^{\frac{k}{\beta}} e^{z}}{\Gamma(\frac{k}{\beta} + \frac{1}{2}) 2^{\frac{k}{\beta}}}. \tag{23}$$

We also have the following obvious inequality.

$$z^{\frac{k-1}{\beta}} e^{-(\frac{1}{2}+\epsilon)z} \leq (\frac{1}{\frac{1}{2}+\epsilon})^{(\frac{k-1}{\beta})} (\frac{k-1}{\beta})^{(\frac{k-1}{\beta})} e^{-\frac{k-1}{\beta}}. \tag{24}$$

From section (12.31) of [50] we get

$$\log \Gamma(\frac{k}{\beta} + \frac{1}{2}) \geq \frac{k}{\beta} \log(\frac{k}{\beta} + \frac{1}{2}) - \frac{k}{\beta} + C. \tag{25}$$

Using (23) and (24), we compute

$$\begin{aligned}
& \Sigma_{k=1}^{\infty} k^2 I_{\frac{k}{\beta}}(z) \\
\leq \quad & \Sigma_{k=1}^{\infty} \frac{\sqrt{\pi} k^2 z^{\frac{k}{\beta}} e^{z}}{\Gamma(\frac{k}{\beta} + \frac{1}{2}) 2^{\frac{k}{\beta}}} = \sqrt{\pi} z^{\frac{1}{\beta}} e^{\frac{3z}{2}} e^{\epsilon z} \Sigma_{k=1}^{\infty} \frac{k^2 z^{\frac{k-1}{\beta}} e^{-(\frac{1}{2}+\epsilon)z}}{\Gamma(\frac{k}{\beta} + \frac{1}{2}) 2^{\frac{k}{\beta}}} \\
\leq \quad & \sqrt{\pi} z^{\frac{1}{\beta}} e^{\frac{3z}{2}} e^{\epsilon z} \Sigma_{k=1}^{\infty} \frac{k^2 (\frac{k-1}{\beta})^{(\frac{k-1}{\beta})} e^{-\frac{k-1}{\beta}} (\frac{2}{1+2\epsilon})^{(\frac{k-1}{\beta})}}{\Gamma(\frac{k}{\beta} + \frac{1}{2}) 2^{\frac{k}{\beta}}} \\
\leq \quad & C z^{\frac{1}{\beta}} e^{\frac{3z}{2}} e^{\epsilon z} \Sigma_{k=1}^{\infty} \frac{(\frac{k}{\beta})^{(\frac{k}{\beta}+1)} e^{-\frac{k}{\beta}}}{\Gamma(\frac{k}{\beta} + \frac{1}{2}) (1+2\epsilon)^{\frac{k}{\beta}}}.
\end{aligned} \tag{26}$$

To estimate the last series above, we take logarithm of the general terms and apply (25) to get

$$\begin{aligned}
&\log[\frac{(\frac{k}{\beta})^{(\frac{k}{\beta}+1)}e^{-\frac{k}{\beta}}}{\Gamma(\frac{k}{\beta}+\frac{1}{2})(1+2\epsilon)^{\frac{k}{\beta}}}] \\
\leq\ & (\frac{k}{\beta}+1)\log(\frac{k}{\beta})-\frac{k}{\beta}-\frac{k}{\beta}\log(\frac{k}{\beta}+\frac{1}{2})+\frac{k}{\beta}+C-\frac{k}{\beta}\log(1+2\epsilon) \\
\leq\ & \log(\frac{k}{\beta})+C-\frac{k}{\beta}\log(1+2\epsilon)\leq C(\epsilon)-\frac{k}{2\beta}\log(1+2\epsilon). \qquad (27)
\end{aligned}$$

Combining (26) and (27), we have

$$\begin{aligned}
&\Sigma_{k=1}^{\infty}k^2 I_{\frac{k}{\beta}}(z) \\
\leq\ & Cz^{\frac{1}{\beta}}e^{\frac{3z}{2}}e^{\epsilon z}\Sigma_{k=1}^{\infty}\frac{(\frac{k}{\beta})^{(\frac{k}{\beta}+1)}e^{-\frac{k}{\beta}}}{\Gamma(\frac{k}{\beta}+\frac{1}{2})(1+2\epsilon)^{\frac{k}{\beta}}}\leq C(\epsilon)z^{\frac{1}{\beta}}e^{\frac{3z}{2}}e^{\epsilon z}\Sigma_{k=1}^{\infty}(1+2\epsilon)^{-\frac{k}{2\beta}} \\
\leq\ & C(\epsilon)z^{\frac{1}{\beta}}e^{\frac{3z}{2}}e^{\epsilon z}.
\end{aligned}$$

□

Now, similar to [17], for any v and $\mu \geq 0$, we define $G_{\mu,v}(r,r',R)$ as

$$G_{\mu,v,h}(r,r',R)=\int_0^{\infty}\int_0^{\infty}(2\pi t)^{1-v}(\frac{rr'}{2t})^h e^{-\lambda^2 t-\frac{R^2}{4t}}J_{\mu}(\lambda r)J_{\mu}(\lambda r')\lambda d\lambda dt.$$

Using formula (17), which is also referable in [49], we have another expression

$$G_{\mu,v,h}(r,r',R)=\pi\int_0^{\infty}(2\pi t)^{-v}(\frac{rr'}{2t})^h e^{-\frac{r^2+r'^2+R^2}{4t}}I_{\mu}(\frac{rr'}{2t})dt. \qquad (28)$$

Lemma 6.2. *Let* $\mu_2>0,\ v>0$. *The following holds:*

- *Suppose* $rr'<\frac{9R^2}{10},\ R>0,\ N\geq\beta\mu_1, h\geq\mu_1-\frac{N}{\beta}$ *(h could be negative). Then*

$$\Sigma_{k\geq\beta\mu_1}k^{\mu_2}|G_{\frac{k}{\beta}-\mu_1,v,h}(r,r',R)|\leq C(v,\mu_1,\mu_2,h)R^{2-2v}.$$

- *Suppose* $r<\frac{4r'}{11},\ r'>0$. *Then*

$$\Sigma_{k\geq\beta\mu_1}k^{\mu_2}|G_{\frac{k}{\beta}-\mu_1,v,h}(r,r',R)|\leq C(v,\mu_1,\mu_2,h)(r')^{2-2v}.$$

Proof. of Lemma 6.2: These estimates are straight forward consequences of Lemma 6.1. By Lemma 6.1 and the assumption that $h - \mu_1 + \frac{N}{\beta} \geq 0$, we have

$$\begin{aligned} & \Sigma_{k\geq N} k^{\mu_2} (\frac{rr^{'}}{2t})^h |G_{\frac{k}{\beta}-\mu_1,v,h}(r, r^{'}, R)| \\ = \ & C\int_0^{+\infty} t^{-v} e^{-\frac{r^2+r^{'2}+R^2}{4t}} (\frac{rr^{'}}{2t})^{h-\mu_1+\frac{N}{\beta}} \Sigma_{k\geq N}^{\infty} k^{\mu_2} I_{\frac{k}{\beta}-\mu_1}(\frac{r\acute{r}}{2t})dt \\ \leq \ & C(v, \mu_1, \mu_2, h, \epsilon)\int_0^{+\infty} t^{-v} e^{-\frac{r^2+r^{'2}+R^2}{4t}} e^{\frac{3r\acute{r}}{4t}} e^{\frac{\epsilon r\acute{r}}{4t}} dt. \end{aligned} \tag{29}$$

We consider the term $e^{-\frac{r^2+r^{'2}+R^2}{4t}} e^{\frac{3r\acute{r}}{4t}} e^{\frac{\epsilon r\acute{r}}{4t}}$ and try to bound it from above in both cases.

When $R > 0$, and $rr^{'} < \frac{9R^2}{10}$, without loss of generality we can assume $R = 1$. Let $\epsilon = 10^{-100}$ and using $rr^{'} < \frac{9}{10}$, we compute

$$\begin{aligned} & e^{-\frac{r^2+r^{'2}+R^2}{4t}} e^{\frac{3r\acute{r}}{4t}} e^{\frac{\epsilon r\acute{r}}{4t}} \\ = \ & e^{-\frac{(r-r^{'})^2}{4t}} e^{-\frac{1-(1+\epsilon)r\acute{r}}{4t}} \leq e^{-\frac{1}{400t}}. \end{aligned}$$

Then in this case we have

$$\begin{aligned} & \Sigma_{k\geq N} k^{\mu_2} |G_{\frac{k}{\beta}-\mu_1,v,h}(r, r^{'}, R)| \\ \leq \ & C(v, \mu_1, \mu_2, h)\int_0^{+\infty} t^{-v} e^{-\frac{1}{400t}} dt \\ \leq \ & C(v, \mu_1, \mu_2, h). \end{aligned}$$

The statement in the first case for all $R > 0$ is thus obtained by rescaling.

In the case when $\acute{r} > 0$ and $r < \frac{4r^{'}}{11}$, we can also assume $\acute{r} = 1$. Then working from (29), using $r < \frac{4}{11}$ and $\epsilon = 10^{-100}$, we have

$$\begin{aligned} & e^{-\frac{r^2+r^{'2}+R^2}{4t}} e^{\frac{3r\acute{r}}{4t}} e^{\frac{\epsilon r\acute{r}}{4t}} \\ \leq \ & e^{-\frac{r^2-(3+\epsilon)r+1}{4t}} \leq e^{-\frac{1}{4\times 10^{10000} t}}, \end{aligned}$$

thus in the second case we simply compute from (29) that

$$\begin{aligned} & \Sigma_{k\geq N} k^{\mu_2} |G_{\frac{k}{\beta}-\mu_1,v,h}(r, r^{'}, R)| \\ \leq \ & C(v, \mu_1, \mu_2, h)\int_0^{+\infty} t^{-v} e^{-\frac{1}{4\times 10^{10000} t}} dt \\ \leq \ & C(v, \mu_1, \mu_2, h). \end{aligned}$$

The proofs in both cases in Lemma 6.2 are completed now. □

7 Local estimates for the heat kernel.

In this section we prove Lemma 7.1,7.2,L7.3, 7.5, 7.6. These lemmas are part of Theorem 1.11.

Lemma 7.1. *For any x and y and $\mathfrak{D} \in \mathfrak{T}$, we have*

$$\begin{cases} \int_0^\infty |\frac{\partial}{\partial\tau}H(x,y,\tau)|d\tau & \leq \quad C|x-y|^{-(m+2)}, \\ \int_0^\infty |\mathfrak{D}H(x,y,\tau)|d\tau & \leq \quad C|x-y|^{-(m+2)}. \end{cases}$$

In particular, we have

$$\begin{cases} \int_0^\infty |\frac{\partial}{\partial\tau}H(0,y,\tau)|d\tau & \leq \quad C|y|^{-(m+2)}, \\ \int_0^\infty |\mathfrak{D}H(0,y,\tau)|d\tau & \leq \quad C|y|^{-(m+2)}. \end{cases}$$

Proof. of Lemma 7.1: We only prove the estimates for the time derivative, the estimate for the spatial derivatives are the same. The estimate $\int_0^\infty |\frac{\partial}{\partial t}H(0,y,\tau)|d\tau \leq C|y|^{-(m+2)}$ has a much shorter proof than the general estimate. To be precise, it suffices to assume $|y| = 1$, assume $r^{'} = 0$ and $\widehat{y} = 0$. Then we have

$$\begin{aligned} & H(0,y,t-\tau) \\ = \quad & \frac{1}{\beta(4\pi t)^{\frac{m}{2}+1}} e^{-\frac{r^2+r^{'2}+R^2}{4t}} [I_0(\frac{rr^{'}}{2t}) + \Sigma_{k\geq 1} 2cosk(\theta-\theta^{'}) I_{\frac{k}{\beta}}(\frac{rr^{'}}{2t})] \\ = \quad & \frac{1}{\beta(4\pi)^{\frac{m}{2}+1}} \frac{1}{t^{\frac{m}{2}+1}} e^{-\frac{1}{4t}}. \end{aligned}$$

Thus

$$\begin{aligned} & \int_0^\infty |\frac{\partial}{\partial t}H(0,y,t-\tau)|d\tau \\ = \quad & \frac{1}{\beta(4\pi)^{\frac{m}{2}+1}} \int_0^\infty |\frac{1}{4t^{\frac{m}{2}+3}} e^{-\frac{1}{4t}} - (\frac{m}{2}+1)\frac{1}{t^{\frac{m}{2}+2}} e^{-\frac{1}{4t}}|d\tau \\ \leq \quad & C(m,\beta). \end{aligned}$$

The general estimate for all x and y is involves more properties of the Bessel functions. We can assume $|x-y| = 1$ and the theorem in general follows from rescaling. Denote $v = \beta(\theta-\theta^{'})$. We still assume that $v \neq \pi$ or $-\pi$ mod $2\beta\pi$ so the last term in (19) vainishes. The conclusion for all v follows from continuity of the estimates with respect to v.

We first compute from (18) that

$$\begin{aligned}
& \frac{\partial}{\partial t}H(x,y,t) \\
= \ & \frac{C}{t^{\frac{m}{2}+1}}[\frac{R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos v)}{4t^2}-(\frac{m}{2}+1)\frac{1}{t}]e^{-\frac{R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos v)}{4t}} \\
& +\frac{C}{t^{\frac{m}{2}+1}}\Sigma_{k\neq 0,\ -\pi<v+2k\beta\pi<\pi}[\frac{R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos(v+2k\beta\pi))}{4t^2} \\
& -(\frac{m}{2}+1)\frac{1}{t}]*e^{-\frac{R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos(v+2k\beta\pi))}{4t}} \\
& +\frac{C}{t^{\frac{m}{2}+1}}[\frac{R^2+(r^{'})^2+r^2}{4t^2}-(\frac{m}{2}+1)\frac{1}{t}]E(\frac{r\acute{r}}{2t},v)*e^{-\frac{R^2+(r^{'})^2+r^2}{4t}} \\
& +\frac{C}{t^{\frac{m}{2}+1}}\frac{r\acute{r}}{2t^2}\frac{\partial E}{\partial z}(\frac{r\acute{r}}{2t},v)e^{-\frac{R^2+(r^{'})^2+r^2}{4t}}.
\end{aligned}$$

It is easy to see from $|x-y|=1$ and (56) we have the following two estimates:

$$R^2+(r^{'}-r)^2+2r\acute{r}\geq\frac{1}{2};$$

$$\begin{aligned}
& R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos(v+2k\beta\pi)) \\
\geq \ & R^2+(r^{'}-r)^2+2r\acute{r}8a \\
\geq \ & R^2+(r^{'}-r)^2+2r\acute{r}(1-\cos v)4a \\
\geq \ & a.
\end{aligned}$$

Then by applying Theorem 10.1 we easily get

$$\frac{\partial}{\partial t}H(x,y,t)|\leq C(1+\frac{1}{t^{\frac{m}{2}+5}})e^{-\frac{a}{1000t}}. \tag{30}$$

Hence

$$\begin{aligned}
& \int_0^\infty|\frac{\partial}{\partial t}H(x,y,t)|dt \\
\leq \ & C\int_0^\infty(1+\frac{1}{t^{\frac{m}{2}+5}})e^{-\frac{a}{1000t}}dt \\
\leq \ & C.
\end{aligned}$$

□

Using similar methods as in the proof of Lemma 7.1, we deduce Lemma 7.2 as follows.

Lemma 7.2. *For any x and y, we have*

$$\int_0^\infty |\nabla_y H(x,y,\tau)| d\tau \leq C|x-y|^{-(m+1)}.$$

Using similar techniques, we also have control on the following integral of third order derivative of the heat kernel.

Lemma 7.3. *For every x and y we have*

$$\int_0^\infty |\nabla_x \frac{\partial}{\partial t} H(x,y,\tau)| d\tau \leq C|y-x|^{-(m+3)}.$$

Proof. of Lemma 7.3: This lemma is a directly corollary of Theorem 10.1. However, we still think it's necessary to give a clear proof here since the lemma so important. It suffices to estimate $\int_0^\infty |\frac{\partial^2}{\partial t \partial r} H(x,y,\tau)| d\tau$, the other derivatives are settled down in the same way. By scaling invariance, we can assume $|x-y| = 1$. Denote $v = \beta(\theta - \theta')$. We still assume that $v \neq \pi$ or $-\pi$ mod $2\beta\pi$ so the last term in (19) vainishes. The conclusion for all v follows from continuity of the estimates with respect to v. The formula says

$$\begin{aligned}
& \frac{\partial^2 H}{\partial t \partial r} \\
= \ & \frac{1}{(4\pi)^{\frac{m}{2}+1}} e^{-\frac{(r-\acute{r})^2 + 2r\acute{r}(1-\cos v) + R^2}{4t}} \{ \frac{1}{t^{\frac{m}{2}+1}} \frac{(r-\acute{r}) + \acute{r}(1-\cos v)}{2t^2} \\
& - \frac{1}{t^{\frac{m}{2}+1}} [\frac{(r-\acute{r})^2 + 2r\acute{r}(1-\cos v) + R^2}{4t^2} - (\frac{m}{2}+1)\frac{1}{t}] \frac{(r-\acute{r}) + \acute{r}(1-\cos v)}{2t} \} \\
& + \frac{1}{(4\pi)^{\frac{m}{2}+1}} \Sigma_{k\neq 0, |v+2k\beta\pi|<\pi} C e^{-\frac{(r-\acute{r})^2 + 2r\acute{r}(1-\cos(v+2k\beta\pi)) + R^2}{4t}} \{ \\
& \frac{1}{t^{\frac{m}{2}+1}} \frac{(r-\acute{r}) + \acute{r}(1-\cos(v+2k\beta\pi))}{2t^2}
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{t^{\frac{m}{2}+1}} [\frac{(r-\acute{r})^2 + 2r\acute{r}(1-\cos(v+2k\beta\pi)) + R^2}{4t^2} - (\frac{m}{2}+1)\frac{1}{t}] \\
& \times \frac{(r-\acute{r}) + \acute{r}(1-\cos(v+2k\beta\pi))}{2t} \} \\
& + \frac{1}{(4\pi)^{\frac{m}{2}+1}} \frac{1}{2\pi\beta} e^{-\frac{r^2+\acute{r}^2+R^2}{4t}} \{ [(\frac{m}{2}+2) \frac{r}{2t^{\frac{m}{2}+3}} - \frac{r}{2t^{\frac{m}{2}+2}} \frac{r^2+\acute{r}^2+R^2}{4t^2}] E(\frac{r\acute{r}}{2t}, v) \\
& + [-(\frac{m}{2}+2) \frac{\acute{r}}{2t^{\frac{m}{2}+3}} + \frac{1}{t^{\frac{m}{2}+1}} \frac{r^2+\acute{r}^2+R^2}{4t^2} * \frac{\acute{r}}{2t} + \frac{1}{t^{\frac{m}{2}+1}} * \frac{r\acute{r}}{2t^2} * \frac{\acute{r}}{2t}] \frac{\partial E(\frac{r\acute{r}}{2t}, v)}{\partial z} \\
& - \frac{(r\acute{r})\acute{r}}{4t^{\frac{m}{2}+4}} \frac{\partial^2 E(\frac{r\acute{r}}{2t}, v)}{\partial z^2} \}.
\end{aligned}$$

Though there are lots of terms, but all the them are actually estimated by a single way. Pick

$$\frac{\acute{r}(1-\cos v)}{t^{\frac{m}{2}+3}}e^{-\frac{(r-\acute{r})^2+R^2}{4t}}e^{-\frac{r\acute{r}}{2t}(1-\cos v)}$$

as an example. First we make the following claim.

Claim 7.4. $\acute{r}(1-\cos v) \leq 4$.

This is easy to prove.
Case 1: Suppose $\acute{r} \leq 2$, it's obvious;
Case 2: Suppose $\acute{r} \geq 2$, since $|x-y| = 1$ we have $r \geq \acute{r} - 1 \geq 1$. Then

$$\acute{r}(1-\cos v) \leq r\acute{r}(1-\cos v) \leq |x-y|^2 = 1.$$

The claim is now proved.

We go on to estimate that

$$\frac{\acute{r}(1-\cos v)}{t^{\frac{m}{2}+3}}e^{-\frac{(r-\acute{r})^2+R^2}{4t}}e^{-\frac{r\acute{r}}{2t}(1-\cos v)} \leq \frac{4}{t^{\frac{m}{2}+3}}e^{-\frac{1}{4t}}.$$

Therefore estimating each term like the above, combining (56), doing the same thing as in Lemma 8.2, we get that

$$|\frac{\partial^2 H}{\partial t \partial r}| \leq C(1+\frac{1}{t^{\frac{m}{2}+5}})e^{-\frac{a}{1000t}},$$

where $a = \frac{\sin^2\frac{\beta\pi}{2}}{4}$. Then

$$\int_0^\infty |\frac{\partial^2 H}{\partial t \partial r}(x,y,t)|dt \leq C.$$

□

It's easy to deduce the following estimate when x is far from the singularity (comparing to $|x-y|$).

Lemma 7.5. *Suppose $P(x) \geq \frac{|x-y|}{100^{100}}$, then for any second order operator $\mathfrak{D} \in \mathfrak{T}$, we have that*

$$\int_0^\infty |\nabla_x \mathfrak{D} H(x,y,\tau)|d\tau \leq C|y-x|^{-(m+3)}.$$

Proof. of Lemma 7.5: It suffices to assume $|x-y| = 1$ and thus from the hypothesis we have that $P(x) \geq \frac{1}{100^{100}}$, which means the r is big. Thus adapting the same proof of Lemma 7.3 , use Theorem 10.1 and $r \geq \frac{1}{100^{100}}$, the proof is completed. In a word, the case when r big is trivial (the real trouble is when r is small). □

Lemma 7.6. *For any x and y, we have*

$$\int_0^\infty |\nabla_x \frac{\partial}{\partial \acute{s}_i} H(x,y,t)|dt \leq C|x-y|^{m+2}$$

and

$$\int_0^\infty |\nabla_x \frac{\partial}{\partial \acute{r}} H(x,y,t)|dt \leq C|x-y|^{m+2}.$$

Notice that the operators $\frac{\partial}{\partial \acute{s}_i}$, $1 \leq i \leq m$ and $\frac{\partial}{\partial \acute{r}}$ are all with respect to the y variable, while ∇_x is with respect to the x variable.

Proof. of Lemma 7.6: This is again a straight forward corollary of Theorem 10.1, just notice that neither $\nabla_x \frac{\partial}{\partial s_i}$ nor $\nabla_x \frac{\partial}{\partial \acute{r}}$ concerns twice or more derivatives on the v (argument) of the $E(z,v)$ in Theorem 10.1.

To be precise, we again assume $|x-y| = 1$ and $v \neq \pi$ or $-\pi$ mod $2\beta\pi$ (so the last term in (19) vainishes). We only show the estimate for $\int_0^\infty |\frac{\partial^2}{\partial r \partial \acute{r}} H(x,y,t)|dt$, the other terms are similar. We compute

$$\begin{aligned}
& \frac{\partial^2}{\partial r \partial \acute{r}} H(x,y,t) \\
= \ & \frac{1}{(4\pi)^{\frac{m}{2}+1}}[\frac{\cos v}{2t^{\frac{m}{2}+2}} + \frac{[(r-\acute{r})+\acute{r}(1-\cos v)][(\acute{r}-r)+r(1-\cos v)]}{4t^{\frac{m}{2}+3}}] \\
\times \ & e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)+R^2}{4t}} + \frac{1}{(4\pi)^{\frac{m}{2}+1}} \Sigma_{k\neq 0, |v+2k\beta\pi|<\pi} [\frac{\cos(v+2k\beta\pi)}{2t^{\frac{m}{2}+2}} \\
& + \frac{[(r-\acute{r})+\acute{r}(1-\cos(v+2k\beta\pi))][(\acute{r}-r)+r(1-\cos(v+2k\beta\pi)))]}{4t^{\frac{m}{2}+3}}] \\
\times \ & e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos(v+2k\beta\pi))+R^2}{4t}} \\
& + \frac{1}{(4\pi)^{\frac{m}{2}+1}} \frac{1}{2\pi\beta} \frac{1}{t^{\frac{m}{2}+1}} [(\frac{r\acute{r}}{4t^2}) E(\frac{r\acute{r}}{2t}, v) - (\frac{r^2}{4t^2}) \frac{\partial}{\partial z} E(\frac{r\acute{r}}{2t}, v) \\
& - (\frac{\acute{r}^2}{4t^2}) \frac{\partial}{\partial z} E(\frac{r\acute{r}}{2t}, v) + (\frac{1}{2t}) \frac{\partial}{\partial z} E(\frac{r\acute{r}}{2t}, v) + (\frac{r\acute{r}}{4t^2}) \frac{\partial^2}{\partial z^2} E(\frac{r\acute{r}}{2t}, v)] e^{-\frac{r^2+\acute{r}^2+R^2}{4t}}.
\end{aligned}$$

The hypothesis $|x-y| = 1$ implies that

$$|(r-\acute{r})+\acute{r}(1-\cos v)| \leq 10, \ |(\acute{r}-r)+r(1-\cos v)| \leq 10; \tag{31}$$

$$\begin{aligned}
& [(r-\acute{r})+\acute{r}(1-\cos(v+2k\beta\pi))][(\acute{r}-r)+r(1-\cos(v+2k\beta\pi)))] \\
\leq\ & (r-\acute{r})^2+(r-\acute{r})(r+\acute{r})(1-\cos(v+2k\beta\pi)) \\
& +r\acute{r}(1-\cos(v+2k\beta\pi))^2 \\
\leq\ & 20+(r-\acute{r})^2+2r\acute{r}(1-\cos(v+2k\beta\pi)).
\end{aligned} \tag{32}$$

Thus, by the idea used in the estimate of $\frac{1}{r}\frac{\partial\widehat{H}}{\partial v}$ in Lemma 9.3, estimate (32), and $|x-y|=1$, it's easy to see that

$$\begin{aligned}
& [(r-\acute{r})+\acute{r}(1-\cos(v+2k\beta\pi))][(\acute{r}-r)+r(1-\cos(v+2k\beta\pi)))] \\
& \times e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos(v+2k\beta\pi))+R^2}{4t}} \\
\leq\ & [20+(r-\acute{r})^2+2r\acute{r}(1-\cos(v+2k\beta\pi))]\times e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos(v+2k\beta\pi))+R^2}{4t}} \\
\leq\ & C(1+t)e^{-\frac{a}{1000t}},
\end{aligned}$$

where a is the one in (56). Applying Theorem 10.1, it's easy to see that

$$|\frac{\partial^2}{\partial r\partial\acute{r}}H(x,y,t)|\leq C(1+\frac{1}{t^{\frac{m}{2}+5}})e^{-\frac{a}{1000t}}.$$

Then we easily get

$$\begin{aligned}
& \int_0^\infty|\frac{\partial^2}{\partial r\partial\acute{r}}H(x,y,t)|dt \\
\leq\ & C\int_0^\infty(1+\frac{1}{t^{\frac{m}{2}+5}})e^{-\frac{a}{1000t}}dt \\
\leq\ & C.
\end{aligned}$$

By rescaling, the proof in the general case is also completed. □

8 Behaviors of the heat kernel near the singular set.

In this section we prove Lemma 8.2. This lemma contains the last two items in Theorem 1.11, and is the only lemma which concerns the exponent $\frac{1}{\beta}-1$.

First of all, the Bessel functions of the second kind satisfies the following recursive relation (see [49]).

$$I_v^{'}(z)+\frac{vI_v(z)}{z}=I_{v-1}(z). \tag{33}$$

Two useful corollaries of the above recursive relation which we will keep applying are

$$\begin{cases} I_v'(z) & \leq & I_{v-1}(z) & \text{when } v \geq 1 \text{ and } z \geq 0; \\ \frac{vI_v(z)}{z} & \leq & I_{v-1}(z) & \text{when } v \geq 1 \text{ and } z \geq 0. \end{cases} \tag{34}$$

Notice that when $v \geq 1$, $\frac{I_v(z)}{z}$ is well defined at $z = 0$.

The following lemma is useful in proving Lemma 8.2.

Lemma 8.1. *Suppose $v \geq 1$ and $z_2 \geq z_1$, we have*

$$|I_v'(z_2) - I_v'(z_1)| \leq \frac{v}{2^v\Gamma(v+1)}|z_2^{v-1} - z_1^{v-1}| + |z_2 - z_1|(1 + \frac{v}{2})I_v(z_2).$$

In particular, if $v < 2$ we have

$$|I_v'(z_2) - I_v'(z_1)| \leq \frac{v}{2^v\Gamma(v+1)}|z_2 - z_1|^{v-1} + |z_2 - z_1|(1 + \frac{v}{2})I_v(z_2).$$

Proof. of Lemma 8.1: Suppose $z_2 \geq z_1$, from the series representation formula in section 3.7 of [49], we compute

$$\begin{aligned} & |I_v'(z_2) - I_v'(z_1)| \\ = \ & \Sigma_{j=0}^{\infty}\frac{(v+2j)z_2^{v+2j-1}}{2^{v+2j}j!(v+j)!} - \Sigma_{j=0}^{\infty}\frac{(v+2j)z_1^{v+2j-1}}{2^{v+2j}j!(v+j)!} \\ = \ & \frac{v}{2^v\Gamma(v+1)}(z_2^{v-1} - z_1^{v-1}) + \Sigma_{j=1}^{\infty}\frac{(v+2j)z_2^{v+2j-1}}{2^{v+2j}j!(v+j)!} - \Sigma_{j=1}^{\infty}\frac{(v+2j)z_1^{v+2j-1}}{2^{v+2j}j!(v+j)!} \\ \leq \ & \frac{v}{2^v\Gamma(v+1)}(z_2^{v-1} - z_1^{v-1}) + (z_2 - z_1)\Sigma_{j=1}^{\infty}\frac{(v+2j)(v+2j-1)z_2^{v+2j-2}}{2^{v+2j}j!(v+j)!}. \end{aligned}$$

Using the simple inequality

$$\frac{(v+2j)(v+2j-1)}{j(v+j)} \leq 4 + 2v, \text{ when } j \geq 1\text{and } j \geq 1,$$

we have

$$\begin{aligned} & \Sigma_{j=1}^{\infty}\frac{(v+2j)(v+2j-1)z_2^{v+2j-2}}{2^{v+2j}j!(v+j)!} \\ \leq \ & (4+2v)\Sigma_{j=1}^{\infty}\frac{z_2^{v+2j-2}}{2^{v+2j}(j-1)!(v+j-1)!} = (1 + \frac{v}{2})I_v(z_2) \end{aligned}$$

Then the proof of the first part is set.

To get the second part it suffices to use $z_2^{v-1} - z_1^{v-1} \leq (z_2 - z_1)^{v-1}$ provided $1 \leq v < 2$. $\square$

Lemma 8.2. *For every second order spatial derivative operator* $\mathfrak{D} \in \mathfrak{T}$*, the following holds. Suppose* $\rho = \min(\frac{1}{\beta} - 1,\ 1)$*,* $|y| = 1$ *and* $|u_1|,\ |u_2| < \frac{1}{8}$*, we have*

$$\int_0^\infty |\mathfrak{D}H(u_1, y, \tau) - \mathfrak{D}H(u_2, y, \tau)| d\tau \leq C|u_1 - u_2|^\rho$$

and

$$\int_0^\infty |\nabla_x H(u_1, y, \tau) - \nabla_x H(u_2, y, \tau)| d\tau \leq C|u_1 - u_2|^\rho.$$

Proof. of Lemma 8.2: Let $u_1 = (r_1, \theta_1, \widehat{u_1})$, $u_2 = (r_2, \theta_2, \widehat{u_2})$, $y = (r', \theta', \widehat{y})$ and $R_1 = |\widehat{y} - \widehat{u_1}|$, $R_2 = |\widehat{y} - \widehat{u_2}|$. We only prove the first inequality in Lemma 8.2, which is a second order estimate. The second inequality is a first order estimate, which is much easier than second order estimates. To prove the first inequality, it suffices to consider the operator $\frac{\partial^2}{\partial r \partial s_i}$, the other operators are similar.

From now on we assume $\frac{1}{\beta} - 1 < 1$, the case when $\frac{1}{\beta} - 1 \geq 1$ is much easier to handle with the following method and we end up with a Lipschitz estimate i.e, with the bound as $C|u_1 - u_2|$.

We split the integral as the following. For all k, write

$$H_k(r, \theta, \acute{r}, \theta', R, t) = \frac{1}{\beta} \frac{1}{(4\pi t)^{\frac{m}{2}+1}} e^{-\frac{r^2 + \acute{r}^2 + R^2}{4t}} I_{\frac{k}{\beta}}\left(\frac{r\acute{r}}{2t}\right) \cos k(\theta - \acute{\theta}).$$

We compute

$$\begin{aligned}
& \int_0^\infty \frac{\partial^2}{\partial r \partial s_i} H_k(u_1, y, t) - \frac{\partial^2}{\partial r \partial s_i} H_k(u_2, y, t) dt \\
= & \int_0^\infty \frac{C}{t^{\frac{m}{2}+3}} [r_1(s_{i,1} - \acute{s}_i) e^{-\frac{r_1^2 + r'^2 + R_1^2}{4t}} I_{\frac{k}{\beta}}(\frac{r_1 r'}{2t}) cos k(\theta_1 - \theta') \\
- & \ r_2(s_{i,2} - \acute{s}_i) e^{-\frac{r_2^2 + r'^2 + R_2^2}{4t}} I_{\frac{k}{\beta}}(\frac{r_2 r'}{2t}) cos k(\theta_2 - \theta')] dt \\
& + \int_0^\infty \frac{C\acute{r}}{t^{\frac{m}{2}+3}} [(s_{i,1} - \acute{s}_i) e^{-\frac{r_1^2 + r'^2 + R_1^2}{4t}} I'_{\frac{k}{\beta}}(\frac{r_1 r'}{2t}) cos k(\theta_1 - \theta') \\
& -(s_{i,2} - \acute{s}_i) e^{-\frac{r_2^2 + r'^2 + R_2^2}{4t}} I'_{\frac{k}{\beta}}(\frac{r_2 r'}{2t}) cos k(\theta_2 - \theta')] dt.
\end{aligned}$$

Thus

$$\int_0^\infty |\frac{\partial^2}{\partial r \partial s_i} H(u_1, y, t) - \frac{\partial^2}{\partial r \partial s_i} H(u_2, y, t)| dt \leq C(|II_1| + |II_2|), \tag{35}$$

where (let $\vartheta_k = 2$ when $k \geq 2$ and $\vartheta_1 = 1$)

$$\begin{aligned} & II_1 \\ = \; & \Sigma_{k=0}^{\infty}\vartheta_k[\int_0^{\infty} |\frac{1}{t^{\frac{m}{2}+3}}[r_1(s_{i,1}-\acute{s}_i)e^{-\frac{r_1^2+r^{'2}+R_1^2}{4t}} I_{\frac{k}{\beta}}(\frac{r_1r^{'}}{2t})cosk(\theta_1-\theta^{'}) \\ - \; & r_2(s_{i,2}-\acute{s}_i)e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}} I_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t})cosk(\theta_2-\theta^{'})]|dt], \end{aligned}$$

and

$$\begin{aligned} & II_2 \\ = \; & \Sigma_{k=0}^{\infty}\int_0^{\infty} |\frac{\acute{r}}{t^{\frac{m}{2}+3}}[(s_{i,1}-\acute{s}_i)e^{-\frac{r_1^2+r^{'2}+R_1^2}{4t}} I^{'}_{\frac{k}{\beta}}(\frac{r_1r^{'}}{2t})cosk(\theta_1-\theta^{'}) \\ & -(s_{i,2}-\acute{s}_i)e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}} I^{'}_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t})cosk(\theta_2-\theta^{'})]|dt. \end{aligned}$$

We only estimate $|II_2|$, and $|II_1|$ can be settled down in the same way and are actually easier to deal with. Let

- $a_{1,k} = cosk(\theta_1-\theta^{'}),\ a_{2,k} = cosk(\theta_2-\theta^{'})$;
- $b_1 = (s_{i,1}-\acute{s}_i)e^{-\frac{r_1^2+r^{'2}+R_1^2}{4t}},\ b_2 = (s_{i,2}-\acute{s}_i)e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}$;
- $c_{1,k} = I^{'}_{\frac{k}{\beta}}(\frac{r_1r^{'}}{2t}),\ c_{2,k} = I^{'}_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t})$.

Using the identity

$$a_{1,k}b_1c_{1,k} - a_{2,k}b_2c_{2,k} = (a_{1,k}-a_{2,k})b_1c_{1,k} + (b_1-b_2)a_{2,k}c_{1,k} + a_{2,k}b_2(c_{1,k}-c_{2,k}), \tag{36}$$

assuming $r_1 \leq r_2$ without loss of generality, we estimate the contributions of $|(a_{1,k}-a_{2,k})b_1c_{1,k}|$, $|(b_1-b_2)a_{2,k}c_{1,k}|$, $|a_{2,k}b_2(c_{1,k}-c_{2,k})|$ to $|II_2|$ independently in the following.

From now on we will repeatedly use the assumption $|u_1|, |u_2|, |y| \leq 1$. These mean that

$$r_1,\ r_2,\ \acute{r},\ \widehat{u_1},\ \widehat{u_2},\ \widehat{y},\ R\ \leq 2. \tag{37}$$

First, on the contribution of $(a_{1,k}-a_{2,k})b_1c_{1,k}$ to $|II_2|$, we compute

$$\begin{aligned} & |(a_{1,k}-a_{2,k})c_{1,k}| \\ = \; & r_1^{\frac{1}{\beta}-1}|cosk(\theta_1-\theta^{'})-cosk(\theta_2-\theta^{'})||[r_1^{1-\frac{1}{\beta}}I^{'}_{\frac{k}{\beta}}(\frac{r_1r^{'}}{2t})]. \end{aligned} \tag{38}$$

Using $r_1 \leq r_2$ and $\sin ku \leq Ck \sin u$, we have

$$\begin{aligned} & r_1^{\frac{1}{\beta}-1} |cosk(\theta_1 - \theta^{'}) - cosk(\theta_2 - \theta^{'})| \\ \leq \quad & Ck\{\sqrt{r_1 r_2 | \sin^2 \frac{|\theta_1 - \theta_2|}{2}}|\}^{\frac{1}{\beta}-1} \leq Ck|u_1 - u_2|^{\frac{1}{\beta}-1}. \end{aligned} \tag{39}$$

By Lemma 8.1, we have for $k \geq 1$ that

$$r_1^{1-\frac{1}{\beta}} I^{'}_{\frac{k}{\beta}}(\frac{r_1 r^{'}}{2t}) \leq r_1^{1-\frac{1}{\beta}} I_{\frac{k}{\beta}-1}(\frac{r_1 r^{'}}{2t}) = (\frac{r^{'}}{2t})^{\frac{1}{\beta}-1} \frac{I_{\frac{k}{\beta}-1}(\frac{r_1 r^{'}}{2t})}{(\frac{r_1 r^{'}}{2t})^{\frac{1}{\beta}-1}}. \tag{40}$$

When $k = 0$ we have

$$r_1^{1-\frac{1}{\beta}} I^{'}_0(\frac{r_1 r^{'}}{2t}) = r_1^{1-\frac{1}{\beta}} I_1(\frac{r_1 r^{'}}{2t}) = (\frac{r^{'}}{2t})^{\frac{1}{\beta}-1} \frac{I_1(\frac{r_1 r^{'}}{2t})}{(\frac{r_1 r^{'}}{2t})^{\frac{1}{\beta}-1}}. \tag{41}$$

Thus for all $k \geq 2$, by (40), (41), (39), and (38), we have

$$\begin{aligned} & |(a_{1,k} - a_{2,k}) c_{1,k}| \\ \leq \quad & Ck(\frac{r^{'}}{2t})^{\frac{1}{\beta}-1} (\frac{r_1 r^{'}}{2t})^{1-\frac{1}{\beta}} I_{\frac{k}{\beta}-1}(\frac{r_1 r^{'}}{2t}) |u_1 - u_2|^{\frac{1}{\beta}-1} \\ \leq \quad & \frac{Ck}{t^{\frac{1}{\beta}-1}} (\frac{r_1 r^{'}}{2t})^{1-\frac{1}{\beta}} I_{\frac{k}{\beta}-1}(\frac{r_1 r^{'}}{2t}) |u_1 - u_2|^{\frac{1}{\beta}-1}. \end{aligned} \tag{42}$$

At this stage, the readers should notice from the assumption $r_1 \leq r_2$ we have that $b_2 \leq b_1$, which is crucial for our trick to work as in the previous steps. Namely, we have to factor $a_1 b_1 c_{1,k} - a_2 b_2 c_{2,k}$ in a right way as in (36). Applying formula (28), Lemma 6.2, and (42), we compute

$$\begin{aligned} & \Sigma_{k=0}^{\infty} \int_0^{\infty} \frac{1}{t^{\frac{m}{2}+3}} |(a_{1,k} - a_{2,k}) c_{1,k}| dt \\ \leq \quad & C|u_1 - u_2|^{\frac{1}{\beta}-1} \Sigma_{k=1}^{\infty} \int_0^{\infty} \frac{k}{t^{\frac{m}{2}+2+\frac{1}{\beta}}} e^{-\frac{r_1^2 + r^{'2} + R_2^2}{4t}} (\frac{r_1 r^{'}}{2t})^{1-\frac{1}{\beta}} I_{\frac{k}{\beta}-1}(\frac{r_1 r^{'}}{2t}) dt \\ = \quad & C|u_1 - u_2|^{\frac{1}{\beta}-1} \Sigma_{k=1}^{\infty} k G_{\frac{k}{\beta}-1, \frac{m}{2}+2+\frac{1}{\beta}, 1-\frac{1}{\beta}}(r_1, r^{'}, R_2) \\ \leq \quad & C|u_1 - u_2|^{\frac{1}{\beta}-1}. \end{aligned} \tag{43}$$

Second, we turn our attention to the contribution of $|(b_1 - b_2) a_{2,k} c_{1,k}|$ to

$|II_2|$. We compute

$$
\begin{aligned}
&|b_1-b_2| = |(s_{i,1}-\acute{s_i})e^{-\frac{r_1^2+r^{'2}+R_1^2}{4t}} - (s_{i,2}-\acute{s_i})e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}| \\
= \;& |(s_{i,1}-\acute{s_i})e^{-\frac{r^{'2}}{4t}}e^{-\frac{R_1^2}{4t}}(e^{-\frac{r_1^2}{4t}}-e^{-\frac{r_2^2}{4t}}) + (s_{i,1}-\acute{s_i})e^{-\frac{r^{'2}}{4t}}e^{-\frac{r_2^2}{4t}}(e^{-\frac{R_1^2}{4t}}-e^{-\frac{R_2^2}{4t}}) \\
&+(s_{i,1}-s_{i,2})e^{-\frac{r^{'2}}{4t}}e^{-\frac{r_2^2+R_2^2}{4t}}|.
\end{aligned}
$$

Then by the mean value theorem and $r_1 \le r_2$ we have

$$|e^{-\frac{r_1^2}{4t}}-e^{-\frac{r_2^2}{4t}}| \le \frac{100|r_1-r_2|}{t}e^{-\frac{r_1^2}{4t}}$$

and

$$|e^{-\frac{R_1^2}{4t}}-e^{-\frac{R_2^2}{4t}}| \le \frac{100|R_1-R_2|}{t}e^{-\frac{R_{1,2}^2}{4t}}$$

for some $R_{1,2}$ between R_1 and R_2. Thus using the same way as we estimate the contribution of $|(a_1-a_2)b_1c_{1,k}|$ to $|II_2|$, we conclude that

$$\Sigma_{k=0}^{\infty}\int_0^{\infty}\frac{1}{t^{\frac{m}{2}+3}}|(b_1-b_2)a_{2,k}c_{1,k}|dt \le C|u_1-u_2|. \qquad (44)$$

Next, we estimate the crucial contribution of $|a_{2,k}b_2(c_{1,k}-c_{2,k})|$ to $|II_2|$.

Using Lemma 8.1 and the inequality $I_1^{'}(z) \le I_0(z)$, recalling that $r_2 \ge r_1$ and $\beta > \frac{1}{2}$, we have

$$
\begin{aligned}
&|c_{1,k}-c_{2,k}| = |I^{'}_{\frac{k}{\beta}}(\frac{r_1r^{'}}{2t}) - I^{'}_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t})| \\
\le \;& \begin{cases}
\frac{\frac{k}{\beta}(\frac{k}{\beta}-1)}{2^{\frac{k}{\beta}}\Gamma(\frac{k}{\beta}+1)}(\frac{r_2r^{'}}{2t})^{\frac{k}{\beta}-2}|\frac{r_2r^{'}}{2t}-\frac{r_1r^{'}}{2t}| + |\frac{r_2r^{'}}{2t}-\frac{r_1r^{'}}{2t}|(1+\frac{k}{2\beta})I_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t}) \\
\qquad \text{when } k \ge 2 \\
\frac{\frac{1}{\beta}}{2^{\frac{1}{\beta}}(\frac{1}{\beta})!}|\frac{r_2r^{'}}{2t}-\frac{r_1r^{'}}{2t}|^{\frac{1}{\beta}-1} + |\frac{r_2r^{'}}{2t}-\frac{r_1r^{'}}{2t}|(1+\frac{1}{2\beta})I_{\frac{1}{\beta}}(\frac{r_2r^{'}}{2t}) \text{ when } k=1 \\
|\frac{r_1r^{'}}{2t}-\frac{r_2r^{'}}{2t}|I_0(\frac{r_2r^{'}}{2t}) \text{ when } k=0.
\end{cases}; \\
= \;& \begin{cases}
\frac{\frac{k}{\beta}(\frac{k}{\beta}-1)}{2^{\frac{k}{\beta}}\Gamma(\frac{k}{\beta}+1)}(\frac{r^{'}}{2t})(\frac{r_2r^{'}}{2t})^{\frac{k}{\beta}-2}|r_2-r_1| + \frac{r^{'}}{2t}|r_2-r_1|(1+\frac{k}{2\beta})I_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t}) \\
\qquad \text{when } k \ge 2 \\
\frac{\frac{1}{\beta}}{2^{\frac{1}{\beta}}(\frac{1}{\beta})!}(\frac{r^{'}}{2t})^{\frac{1}{\beta}-1}|r_2-r_1|^{\frac{1}{\beta}-1} + |r_2-r_1|\frac{r^{'}}{2t}(1+\frac{1}{2\beta})I_{\frac{1}{\beta}}(\frac{r_2r^{'}}{2t}) \text{ when } k=1 \\
\frac{r^{'}}{2t}|r_2-r_1|I_0(\frac{r_2r^{'}}{2t}) \text{ when } k=0.
\end{cases};
\end{aligned}
$$

When $k = 0$ we implicitly use $I_{-1}(x) = I_1(x)$, thanks to the properties of Bessel functions.

We estimate the contribution of $|a_{2,k}b_2(c_{1,k} - c_{2,k})|$ to $|II_2|$ as

$$\begin{aligned}
& \Sigma_{k\geq 2}\int_0^\infty \frac{1}{t^{\frac{m}{2}+3}}|c_{1,k} - c_{2,k}|a_{2,k}b_2 dt \\
& + \int_0^\infty \frac{1}{t^{\frac{m}{2}+3}}|c_{1,1} - c_{2,1}|a_{2,1}b_2 dt + \int_0^\infty \frac{1}{t^{\frac{m}{2}+3}}|c_{1,0} - c_{2,0}|a_{2,0}b_2 dt \\
\leq \; & N_1 + N_2 + N_3 + N_4 + N_5, \qquad (45)
\end{aligned}$$

where

$$\begin{aligned}
N_1 &= \Sigma_{k\geq 2}\int_0^\infty \frac{\frac{k}{\beta}(\frac{k}{\beta} - 1)}{2^{\frac{k}{\beta}}\Gamma(\frac{k}{\beta}+1)}(\frac{r^{'}}{2t})(\frac{r_2r^{'}}{2t})^{\frac{k}{\beta}-2}|r_2 - r_1|\frac{|s_{i,2} - \acute{s_i}|}{t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}dt, \\
N_2 &= \Sigma_{k\geq 2}\int_0^\infty (1 + \frac{k}{2\beta})(\frac{r^{'}}{2t})|r_2 - r_1|I_{\frac{k}{\beta}}(\frac{r_2r^{'}}{2t})\frac{|s_{i,2} - \acute{s_i}|}{t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}dt, \\
N_3 &= \int_0^\infty \frac{\frac{1}{\beta}}{2^{\frac{1}{\beta}}(\frac{1}{\beta})!}(\frac{r^{'}}{2t})^{\frac{1}{\beta}-1}|r_2 - r_1|^{\frac{1}{\beta}-1}\frac{|s_{i,2} - \acute{s_i}|}{t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}dt, \\
N_4 &= \int_0^\infty (1 + \frac{1}{2\beta})(\frac{r^{'}}{2t})|r_2 - r_1|I_{\frac{1}{\beta}}(\frac{r_2r^{'}}{2t})\frac{|s_{i,2} - \acute{s_i}|}{t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}, \\
N_5 &= \int_0^\infty (\frac{r^{'}}{2t})|r_2 - r_1|I_0(\frac{r_2r^{'}}{2t})\frac{|s_{i,2} - \acute{s_i}|}{2t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}dt.
\end{aligned}$$

Before estimating term by term, we should deal with the term $e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}$ first.

Claim 8.3. $e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}} \leq e^{-\frac{1}{8t}}$.

The see why the claim holds, just notice that we have $|u_1|, |u_2| \leq \frac{1}{8}$, $r^{'2} + |\widehat{y}|^2 = |y|^2 = 1$, then we get

$$\begin{aligned}
& r^{'2} + R_2^2 = r^{'2} + (\widehat{y} - \widehat{u}_2)^2 \\
\geq \; & 1 - [|\widehat{y}|^2 - ||\widehat{y}| - |\widehat{u_2}||^2] \\
= \; & 1 - |\widehat{u_2}|(|\widehat{y}| + |\widehat{u_2}|) \geq 1 - \frac{1}{8} \times \frac{9}{8} > \frac{1}{2}.
\end{aligned}$$

Therefore $e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}} \leq e^{-\frac{1}{8t}}$.

Now we estimate term by term. For N_1, we compute

$$\begin{aligned}
& N_1 \\
= \; & \Sigma_{k\geq 2}\int_0^\infty \frac{\frac{k}{\beta}(\frac{k}{\beta}-1)}{2^{\frac{k}{\beta}}\Gamma(\frac{k}{\beta}+1)}(\frac{r^{'}}{2t})(\frac{r_2r^{'}}{2t})^{\frac{k}{\beta}-2}|r_2-r_1|\frac{|s_{i,2}-\acute{s}_i|}{t^{\frac{m}{2}+3}}e^{-\frac{r_2^2+r^{'2}+R_2^2}{4t}}dt \\
\leq \; & |u_1-u_2|\Sigma_{k\geq 2}\frac{\frac{k}{\beta}(\frac{k}{\beta}-1)}{2^{\frac{2k}{\beta}-2}\Gamma(\frac{k}{\beta}+1)8^{\frac{k}{\beta}-2}}\int_0^\infty \frac{1}{t^{\frac{m}{2}+2+\frac{k}{\beta}}}e^{-\frac{1}{8t}}dt.
\end{aligned}$$

Notice that by the definition of the *Gamma*-function we have

$$\int_0^\infty \frac{1}{t^{\frac{m}{2}+2+\frac{k}{\beta}}}e^{-\frac{1}{8t}}dt = 8^{\frac{m}{2}+\frac{k}{\beta}+1}\Gamma(\frac{m}{2}+\frac{k}{\beta}+1).$$

Then we have

$$\begin{aligned}
& N_1 \\
\leq \; & |u_1-u_2|\Sigma_{k\geq 2}\frac{\frac{k}{\beta}(\frac{k}{\beta}-1)(8^{\frac{m}{2}+\frac{k}{\beta}+1})\Gamma(\frac{m}{2}+\frac{k}{\beta}+1)}{2^{\frac{2k}{\beta}-1}\Gamma(\frac{k}{\beta}+1)8^{\frac{k}{\beta}-2}} \\
\leq \; & |u_1-u_2|\Sigma_{k\geq 2}\frac{(\frac{k}{\beta})^2(8^{\frac{m}{2}+4})(\frac{k}{\beta}+m)^m}{4^{\frac{k}{\beta}}} \\
\leq \; & C|u_1-u_2|.
\end{aligned}$$

To deduce the second inequality above we used the fact that $\frac{\Gamma(\frac{m}{2}+\frac{k}{\beta}+1)}{\Gamma(\frac{k}{\beta}+1)} \leq (\frac{k}{\beta}+m)^m$. Thus we finished the estimate on N_1.

For N_2, from formula (28), and Corollary 6.2, taking $v=8$, $\mu_1=\frac{1}{8}$ and $\mu_2=1$ in Corollary 6.2, we have that

$$\begin{aligned}
& N_2 \\
\leq \; & C|u_2-u_1|\Sigma_{k\geq 2}(1+\frac{k}{2\beta})G_{\frac{k}{\beta},\frac{m}{2}+4,0}(r_2,r^{'},R_2) \\
\leq \; & C|u_2-u_1|.
\end{aligned}$$

For N_3, actually it's our crucial term since it concerns the Hölder continuity exponent $\frac{1}{\beta}-1$. Nevertheless, from our established results, N_3 can be estimated easily as

$$\begin{aligned}
& N_3 \\
\leq \; & C|u_1-u_2|^{\frac{1}{\beta}-1}\int_0^\infty \frac{1}{t^{\frac{m}{2}+2+\frac{1}{\beta}}}e^{-\frac{1}{8t}}dt \\
\leq \; & C|u_1-u_2|^{\frac{1}{\beta}-1}.
\end{aligned}$$

Using the same (and easier) way as we estimated N_2, we have that

$$N_4 + N_5 \leq C|u_1 - u_2|,$$

where we applied Claim 8.3. Since $|u_1 - u_2| \leq |u_1 - u_2|^{\frac{1}{\beta}-1} \leq 1$, using the decomposition (45) and the above estimates for $(N_i, 0 \leq i \leq 5)$, we estimate the contribution of $|c_{1,k} - c_{2,k}|a_{2,k}b_2$ to $|II_2|$ as

$$\Sigma_{k=0}^{\infty} \int_0^{\infty} \frac{1}{t^{\frac{m}{2}+3}} |c_{1,k} - c_{2,k}|a_{2,k}b_2 dt \leq C|u_1 - u_2|^{\frac{1}{\beta}-1}. \tag{46}$$

Thus now we are ready to make the final conclusion on $|II_2|$. Assume $\frac{1}{\beta} - 1 < 1$, using (43), (44), and (46) on the contributions of $|(a_1 - a_2)b_1 c_{1,k}|$, $|(b_1 - b_2)a_2 c_{1,k}|$, $|a_2 b_2 (c_{1,k} - c_{2,k})|$ to $|II_2|$, we have

$$|II_2| \leq C|u_1 - u_2|^{\frac{1}{\beta}-1}. \tag{47}$$

Regarding $|II_1|$, the Hölder continuity exponent $\frac{1}{\beta} - 1$ doesn't appear in it. Actually employing the same method as we estimate $|II_2|$ (in a much easier way), the following estimate holds.

$$|II_1| \leq C|u_1 - u_2|. \tag{48}$$

Thus by (35), (48), and (47), the proof of the lemma is completed. □

9 Asymptotic behaviors of the heat kernel.

In this section we prove Lemma 9.1 and proposition 9.2, which are two items in Theorem 1.11. They rely on the asymptotic estimates for the derivatives of the heat kernel in Lemma 9.4, 9.3, 9.5, which are implied by Theorem 10.1.

From now on, in both section 9 and 10 we write $v = \beta(\theta - \acute{\theta})$. Notice that the domain of v is $[-\beta\pi, \beta\pi]$.

Lemma 9.1. *Suppose* $t = 1$, $x \in A_{\frac{1}{\sqrt{s}}}$, $y \in A_{\frac{10}{\sqrt{s}}}$, $0 < s \leq \frac{1}{10000}$, *then the following estimate holds*

$$|\nabla_x H(x, y, 1)| \leq Ce^{-\frac{1}{s}}.$$

Proof. of Lemma 9.1: This is a direct consequence of the lemmas in section 9. We only consider the estimate for $\frac{1}{r}\frac{\partial}{\partial\theta}H(x,y,1)$, the other directional derivatives are similar. From the hypothesis

$$r^2+|\widehat{x}|^2=\frac{1}{s},\ \acute{r}^2+|\widehat{y}|^2=\frac{100}{s},$$

it's easy to deduce that

$$(r-r^{'})^2+R^2\geq\frac{20}{s}.$$

Then from Lemma 9.4 and 9.3, it's straightforward to see that

$$|\frac{1}{r}\frac{\partial}{\partial\theta}H(x,y,1)|\leq Ce^{-\frac{1}{s}}.$$

$\square$

Let $a=\frac{\sin^2(\frac{\beta\pi}{2})}{4}$. In the following discussions, we will frequently use the inequality

$$1+x^p\leq C(1+x^q)\text{ if }q\geq p\text{ and }0\leq p\leq q\leq 10, \tag{49}$$

and the inequality

$$x^pe^{-x^q}\leq C_{p,q},\ \text{if }q>0. \tag{50}$$

Proposition 9.2. *There exists a constant C with the following property. For any $x\in\mathbb{R}^2\times\mathbb{R}^m$, any second order spatial differential operator $\mathfrak{D}\in\mathfrak{T}$, and any α such that $0\leq\alpha\leq 1$, the following estimates on the heat kernel hold.*

$$\int_{\mathbb{R}^2\times\mathbb{R}^m}|\sup_{1\leq t\leq 2}\frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)||x-y|^{\alpha}dy\leq C;$$

$$\int_{\mathbb{R}^2\times\mathbb{R}^m}|\sup_{1\leq t\leq 2}\frac{\partial^2}{\partial t^2}H(x,y,t)||x-y|^{\alpha}dy\leq C.$$

Proof. of Proposition 9.2: We still write $v=\beta(\theta-\acute{\theta})$. Since

$$H=\widehat{H}\times H_E,\ H_E=\frac{1}{(4\pi t)^{\frac{m}{2}}}e^{-\frac{R^2}{4t}},\ R=|\widehat{x}-\widehat{y}|, \tag{51}$$

we obviously have

$$\begin{aligned}&|\sup_{1\leq t\leq 2}\frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)|\\ \leq\ &(\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial\widehat{H}}{\partial\theta}|+\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial^2\widehat{H}}{\partial t\partial\theta}|+\sup_{1\leq t\leq 2}|\frac{\partial\widehat{H}}{\partial r}|+\sup_{1\leq t\leq 2}|\frac{\partial^2\widehat{H}}{\partial t\partial r}|+\sup_{1\leq t\leq 2}|\frac{\partial\widehat{H}}{\partial t}|)\\ &\times(\sup_{1\leq t\leq 2}|\frac{\partial H_E}{\partial t}|+\Sigma_{i=1}^m(\sup_{1\leq t\leq 2}|\frac{\partial H_E}{\partial s_i}|+\sup_{1\leq t\leq 2}|\frac{\partial^2 H_E}{\partial t\partial s_i}|)).\end{aligned}$$

From (51) we see that

$$(\sup_{1\leq t\leq 2}|\frac{\partial H_E}{\partial t}|+\Sigma_{i=1}^m(\sup_{1\leq t\leq 2}|\frac{\partial H_E}{\partial s_i}|+\sup_{1\leq t\leq 2}|\frac{\partial^2 H_E}{\partial t\partial s_i}|)\leq Ce^{-\frac{R^2}{16}}. \tag{52}$$

Without loss of generality we can assume $\theta=0$. Thus from Lemma 9.4, 9.3, and 9.5, we have

$$\begin{aligned}
&(\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial \widehat{H}}{\partial \theta}|+\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial^2 \widehat{H}}{\partial t\partial \theta}|+\sup_{1\leq t\leq 2}|\frac{\partial \widehat{H}}{\partial r}| \\
&+\sup_{1\leq t\leq 2}|\frac{\partial^2 \widehat{H}}{\partial t\partial r}|+\sup_{1\leq t\leq 2}|\frac{\partial \widehat{H}}{\partial t}|) \\
\leq\ & C\underline{Z},
\end{aligned}$$

where

$$\begin{aligned}
& \underline{Z} \\
=\ & (1+\frac{1}{r}|r\acute{r}\sin v|)e^{-\frac{(r-\acute{r})^2}{16}}e^{-\frac{r\acute{r}[1-\cos v]}{8}} \\
& +e^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}+e^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Thus

$$|\sup_{1\leq t\leq 2}\frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)|\leq C\underline{Z}e^{-\frac{R^2}{16}}.$$

Then combining 52, we estimate

$$\begin{aligned}
&\int_{\mathbb{R}^2\times\mathbb{R}^m}|\sup_{1\leq t\leq 2}\frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)||x-y|^\alpha dy \\
\leq\ & C\int_{-\beta\pi}^{\beta\pi}\int_0^\infty\int_{R^m}\underline{Z}e^{-\frac{R^2}{16}}(|r-\acute{r}|^\alpha+r\acute{r}[1-\cos v]+R^\alpha)\acute{r}d\acute{r}dvd\widehat{y} \\
=\ & C\int_{-\beta\pi}^{\beta\pi}\int_0^\infty\underline{Z}(|r-\acute{r}|^\alpha+r\acute{r}[1-\cos v])\acute{r}d\acute{r}dv\int_{R^m}e^{-\frac{R^2}{16}}d\widehat{y} \\
& +C\int_{-\beta\pi}^{\beta\pi}\int_0^\infty\underline{Z}\acute{r}d\acute{r}dv\int_{R^m}R^\alpha e^{-\frac{R^2}{16}}d\widehat{y}.
\end{aligned}$$

Since it's obvious that

$$\int_{R^m}R^\alpha e^{-\frac{R^2}{16}}d\widehat{y}+\int_{R^m}e^{-\frac{R^2}{16}}d\widehat{y}\leq C,$$

we have

$$\begin{aligned}
& \int_{\mathbb{R}^2\times\mathbb{R}^m} |\sup_{1\le t\le 2}\frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)||x-y|^\alpha dy \\
\le\ & \int_{-\beta\pi}^{\beta\pi}\int_0^\infty \underline{Z}(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos\theta^{'}])^{\frac{\alpha}{2}})\acute{r}d\acute{r}dv \\
\le\ & U_1+U_2+U_3+U_4.
\end{aligned}$$

The four terms in the decomposition are

$$\begin{aligned}
U_1 &= \int_{-\beta\pi}^{\beta\pi}\int_0^\infty e^{-\frac{(r-\acute{r})^2}{16}}e^{-\frac{r\acute{r}[1-\cos v]}{8}}|r\acute{r}\sin v|(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos v])^{\frac{\alpha}{2}}) \\
&\quad \times\frac{\acute{r}}{r}d\acute{r}dv, \\
U_2 &= \int_{-\beta\pi}^{\beta\pi}\int_0^\infty e^{-\frac{(r-\acute{r})^2}{16}}e^{-\frac{r\acute{r}[1-\cos v]}{8}}(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos v])^{\frac{\alpha}{2}})\acute{r}d\acute{r}dv, \\
U_3 &= \int_{-\beta\pi}^{\beta\pi}\int_0^\infty e^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos v])^{\frac{\alpha}{2}})\acute{r}d\acute{r}dv, \\
U_4 &= \int_{-\beta\pi}^{\beta\pi}\int_0^\infty e^{-\frac{r^2+\acute{r}^2}{16}}(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos v])^{\frac{\alpha}{2}})\acute{r}d\acute{r}dv.
\end{aligned}$$

We first estimate the most critical term U_1. Using (50) we have that

$$e^{-\frac{(r-\acute{r})^2}{32}}e^{-\frac{r\acute{r}[1-\cos v]}{16}}(1+|r-\acute{r}|^\alpha+(r\acute{r}[1-\cos v])^{\frac{\alpha}{2}})\le C. \tag{53}$$

Then

$$U_1\le C\int_{-\beta\pi}^{\beta\pi}\int_0^\infty e^{-\frac{(r-\acute{r})^2}{32}}e^{-\frac{r\acute{r}[1-\cos v]}{16}}|r\acute{r}\sin v|\frac{\acute{r}}{r}d\acute{r}dv. \tag{54}$$

Notice that we have

$$\begin{aligned}
\textstyle\int_{-\beta\pi}^{\beta\pi}|r\acute{r}\sin v|e^{-\frac{r\acute{r}[1-\cos v]}{16}}dv &= 2\textstyle\int_0^{\beta\pi}(r\acute{r}\sin v)e^{-\frac{r\acute{r}[1-\cos v]}{16}}dv \\
&= 32[1-e^{-\frac{r\acute{r}[1-\cos\beta\pi]}{16}}].
\end{aligned}$$

To bound U_1, using the above identity we compute

$$\begin{aligned}
U_1 &\le C\int_0^\infty\frac{\acute{r}}{r}e^{-\frac{(r-\acute{r})^2}{32}}d\acute{r}\int_{-\beta\pi}^{\beta\pi}e^{-\frac{r\acute{r}[1-\cos v]}{16}}|r\acute{r}\sin v|dv \\
&\le C\int_0^\infty\frac{\acute{r}}{r}e^{-\frac{(r-\acute{r})^2}{32}}[1-e^{-\frac{r\acute{r}[1-\cos\beta\pi]}{16}}]d\acute{r}.
\end{aligned} \tag{55}$$

When $r \geq 1$, from (56) we have

$$\begin{aligned} U_1 &\leq C\Big[\int_0^{\infty} \frac{\acute{r}-r}{r} e^{-\frac{(r-\acute{r})^2}{32}} d\acute{r} + \int_0^{\infty} e^{-\frac{(r-\acute{r})^2}{32}} d\acute{r}\Big] \\ &\leq C. \end{aligned}$$

When $r \leq 1$, using the inequality

$$[1 - e^{-\frac{r\acute{r}[1-\cos\beta\pi]}{16}}] \leq Cr \cdot \acute{r}$$

and the inequality

$$\acute{r}^2 \leq (\acute{r}-r)^2 + 2|\acute{r}-r| + 1,$$

we obtain from (56) that

$$\begin{aligned} U_1 &\leq C\int_0^{\infty} \frac{\acute{r}}{r} e^{-\frac{(r-\acute{r})^2}{32}} [1 - e^{-\frac{r\acute{r}[1-\cos\beta\pi]}{16}}] d\acute{r} \leq C\int_0^{\infty} \acute{r}^2 e^{-\frac{(r-\acute{r})^2}{32}} d\acute{r} \\ &\leq C\int_0^{\infty} ((\acute{r}-r)^2 + 2|\acute{r}-r| + 1) e^{-\frac{(r-\acute{r})^2}{32}} d\acute{r} \\ &\leq C. \end{aligned}$$

Thus, U_1 is bounded uniformly. Using (50), it's easy to see that

$$U_2,\ U_3,\ U_4 \leq C.$$

Therefore, every term is controlled so the estimate of

$$\int_{\mathbb{R}^2\times\mathbb{R}^m} |\sup_{1\leq t\leq 2} \frac{\partial}{\partial t}\mathfrak{D}H(x,y,t)||x-y|^{\alpha} dy$$

follows.

The estimate of $\int_{\mathbb{R}^2\times\mathbb{R}^m} |\sup_{1\leq t\leq 2} \frac{\partial^2}{\partial t^2} H(x,y,t)||x-y|^{\alpha} dy$ is a directly consequence of Lemma 9.5 and similar arguments as above. $\square$

Lemma 9.3. *We have*

$$\begin{aligned} &\sup_{1\leq t\leq 2} |\frac{1}{r}\frac{\partial \widehat{H}}{\partial \theta}| + \sup_{1\leq t\leq 2} |\frac{1}{r}\frac{\partial^2 \widehat{H}}{\partial t\partial \theta}| \\ \leq\ & \frac{C}{r} e^{-\frac{(r-\acute{r})^2}{16}} e^{-\frac{r\acute{r}[1-\cos\beta(\theta-\theta')]}{8}} |r\acute{r}\sin\beta(\theta-\theta')| + Ce^{-\frac{(r-\acute{r})^2}{16}} e^{-ar\acute{r}} \\ &+ Ce^{-\frac{r^2+\acute{r}^2}{16}}. \end{aligned}$$

Proof. of Lemma 9.3: We still write $v = \beta(\theta - \theta')$. We further assume that $v \neq \pi$ or $-\pi \mod 2\beta\pi$ so the last term in (19) vainishes. The conclusion for all v follows from continuity. There is an interesting point hiding here: though $E(v, z)$ is not continuous with respect to v, the estimates we get by applying the bounds on $E(v, z)$ still works for all v, as the estimates are all continuous.

We compute

$$\begin{aligned}
& \frac{1}{\beta r}\frac{\partial \widehat{H}}{\partial \theta} = \frac{1}{r}\frac{\partial \widehat{H}}{\partial v} \\
= \ & \frac{1}{r}\frac{1}{4\pi t}e^{-\frac{r^2+\acute{r}^2}{4t}}[-\Sigma_{k,|v+2k\beta\pi|<\pi}\frac{r\acute{r}}{2t}\sin(v+2k\beta\pi)e^{\frac{r\acute{r}}{2t}\cos(v+2k\beta\pi)} + \frac{1}{2\pi\beta}\frac{\partial E(\frac{r\acute{r}}{2t}, v)}{\partial v}] \\
= \ & \frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}e^{-\frac{r^2+\acute{r}^2}{4t}}\frac{\partial E(\frac{r\acute{r}}{2t}, v)}{\partial v} - \frac{1}{r}\frac{1}{4\pi t}e^{-\frac{(r-\acute{r})^2}{4t}}\frac{r\acute{r}}{2t}(\sin v)e^{-\frac{r\acute{r}}{2t}(1-\cos v)} \\
& -\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}\frac{1}{r}\frac{1}{4\pi t}e^{-\frac{(r-\acute{r})^2}{4t}}\frac{r\acute{r}}{2t}[\sin(v+2k\beta\pi)]e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}.
\end{aligned}$$

Next, note that when $k \neq 0$, we have

$$1 - \cos(v + 2k\beta\pi) \geq 2\sin^2(\frac{\beta\pi}{2}) = 8a. \tag{56}$$

Combining the estimates in Theorem 10.1, we have

$$\begin{aligned}
& \sup_{1\leq t\leq 2}|\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}\frac{1}{r}\frac{1}{4\pi t}e^{-\frac{(r-\acute{r})^2}{4t}}\frac{r\acute{r}}{2t}[\sin(v+2k\beta\pi)]e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}| \\
& \quad \leq \frac{C}{r}(r\acute{r})e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \leq Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}.
\end{aligned} \tag{57}$$

Inequality (57) is proved as the following.

1. When $r \geq 1$, obviously we have
$$\frac{C}{r}(r\acute{r})e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \leq Ce^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}};$$
2. when $r \leq 1,\ \acute{r} \leq 2$, we estimate
$$C\acute{r}e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \leq Ce^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}};$$
3. when $r \leq 1,\ \acute{r} \geq 2$, we estimate
$$C\acute{r}e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \leq C(\acute{r}-r)e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \leq Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-2ar\acute{r}}.$$

By Theorem 10.1 we immediately get

$$\begin{aligned}\sup_{1\le t\le 2}\left|\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}e^{-\frac{r^2+\acute{r}^2}{4t}}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v}\right| &\le C\acute{r}\sup_{1\le t\le 2}e^{-\frac{r^2+\acute{r}^2}{8}}\left|\frac{1}{(\frac{r\acute{r}}{2t})}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v}\right|\\ &\le Ce^{-\frac{r^2+\acute{r}^2}{16}}.\end{aligned}$$

Thus, we conclude

$$\begin{aligned}&\sup_{1\le t\le 2}\left|\frac{1}{r}\frac{\partial\widehat{H}}{\partial v}\right|\\ \le\ &\frac{C}{r}e^{-\frac{(r-\acute{r})^2}{8}}e^{-\frac{r\acute{r}[1-\cos v]}{4}}|r\acute{r}\sin v|+Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}\\ &+Ce^{-\frac{r^2+\acute{r}^2}{16}}.\end{aligned}\tag{58}$$

Continue differentiating (56) with respect to t we get

$$\begin{aligned}&\frac{1}{\beta r}\frac{\partial^2\widehat{H}}{\partial t\partial\theta}=\frac{1}{r}\frac{\partial^2\widehat{H}}{\partial t\partial v}\\ =\ &e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)}{4t}}\{\frac{1}{r}\frac{r\acute{r}\sin v}{4\pi t^3}-\frac{1}{r}\frac{r\acute{r}\sin v}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}-\frac{1}{r}\frac{r\acute{r}\sin v}{8\pi t^2}\frac{r\acute{r}}{2t^2}(1-\cos v)\}\\ &+\Sigma_{k\ne 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{4\pi t^3}\\ &-\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}-\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{8\pi t^2}\frac{r\acute{r}}{2t^2}[1-\cos(v+2k\beta\pi)]\}\\ &+e^{-\frac{r^2+\acute{r}^2}{4t}}\{-\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t^2}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v}+\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}(\frac{r^2+\acute{r}^2}{4t^2})\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v}\\ &+\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}\frac{\partial^2 E(\frac{r\acute{r}}{2t},v)}{\partial z\partial v}(-\frac{r\acute{r}}{2t^2})\}.\end{aligned}$$

Then, by the same idea as in the estimate of $\sup_{1\le t\le 2}|\frac{1}{r}\frac{\partial\widehat{H}}{\partial v}|$, using (56), Theorem 10.1, and Cauchy-Schwarz inequalities, we directly get

$$\begin{aligned}&\sup_{1\le t\le 2}|\Sigma_{k\ne 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{4\pi t^3}\\ &-\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}-\frac{1}{r}\frac{r\acute{r}\sin(v+2k\beta\pi)}{8\pi t^2}\frac{r\acute{r}}{2t^2}[1-\cos(v+2k\beta\pi)]\}|\\ \le\ &Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}\end{aligned}$$

and

$$\begin{aligned}
& \sup_{1\leq t\leq 2} |e^{-\frac{r^2+\acute{r}^2}{4t}}\{-\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t^2}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v}+\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t^2}(\frac{r^2+\acute{r}^2}{4t})\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial v} \\
& +\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}\frac{\partial^2 E(\frac{r\acute{r}}{2t},v)}{\partial z\partial v}(-\frac{r\acute{r}}{2t^2})\}| \\
\leq\ & C(1+r^3+\acute{r}^3)e^{-\frac{r^2+\acute{r}^2}{8}}, \\
\leq\ & Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Thus, the estimate of $\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial^2\widehat{H}}{\partial t\partial v}|$ follows.

The conclusion of Lemma 9.3 directly follows from (58) and (59). ☐

The other lemmas in this section are proved under the same ideas as in the proof of Lemma 9.3. For the sake of clarity, we still give detailed proof here, since these lemmas are crucial to the global estimates of the derivatives of the heat kernel and their conclusions are not the same.

Lemma 9.4. *We have*

$$\begin{aligned}
\sup_{1\leq t\leq 2}|\frac{\partial\widehat{H}}{\partial r}|+\sup_{1\leq t\leq 2}|\frac{\partial^2\widehat{H}}{\partial t\partial r}| \ \leq\ & Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-\frac{r\acute{r}[1-\cos\beta(\theta-\theta^{'})]}{8}}+Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}} \\
& +Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Proof. of Lemma 9.4: Denote $v=\beta(\theta-\theta^{'})$. We further assume that $v\neq\pi$ or $-\pi$ mod $2\beta\pi$ so the last term in (19) vainishes. The conclusion for all v follows from continuity of the estimates with respect to v.
First we compute

$$\begin{aligned}
\frac{\partial\widehat{H}}{\partial r} \ =\ & e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)}{4t}}\{-\frac{(r-\acute{r})}{8\pi t^2}-\frac{1}{r}\frac{r\acute{r}(1-\cos v)}{8\pi t^2}\} \\
& +\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2}{4t}}e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}\{-\frac{(r-\acute{r})}{8\pi t^2} \\
& -\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^2}\} \\
& +e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}\frac{r\acute{r}}{2t}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{4\pi t}\frac{r}{2t}E(\frac{r\acute{r}}{2t},v)\}.
\end{aligned}$$

Then by the same idea as in the estimate of $\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial\widehat{H}}{\partial v}|$, using (56), Theorem 10.1, (49), and (50), we get

$$\begin{aligned}
&\sup_{1\leq t\leq 2}|\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2}{4t}}e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}\{-\frac{(r-\acute{r})}{8\pi t^2}\\
&-\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^2}\}|\\
\leq\;& C(|r-\acute{r}|+\acute{r})e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}}\\
\leq\;& Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}.
\end{aligned}$$

Then using the simple estimate $e^{-\frac{r^2+\acute{r}^2}{16}}(1+r^2)\leq C$ we have

$$\begin{aligned}
&\sup_{1\leq t\leq 2}|e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{1}{r}\frac{1}{4\pi t}\frac{r\acute{r}}{2t}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{4\pi t}\frac{r}{2t}E(\frac{r\acute{r}}{2t},v)\}|\\
\leq\;& Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Thus the estimate for $\sup_{1\leq t\leq 2}|\frac{\partial\widehat{H}}{\partial r}|$ follows.

Continue differentiating (59) with respect to t we get

$$\begin{aligned}
&\frac{\partial^2\widehat{H}}{\partial t\partial r}\\
=\;& e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)}{4t}}\{\frac{(r-\acute{r})}{4\pi t^3}-\frac{(r-\acute{r})}{16\pi t^4}[r\acute{r}(1-\cos v)]-\frac{(r-\acute{r})^3}{32\pi t^4}\\
&+\frac{1}{r}\frac{r\acute{r}(1-\cos v)}{4\pi t^3}-\frac{1}{r}\frac{r\acute{r}(1-\cos v)}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}-\frac{1}{r}\frac{[r\acute{r}(1-\cos v)]^2}{16\pi t^4}\}\\
+\;& \Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{(r-\acute{r})}{4\pi t^3}\\
&-\frac{(r-\acute{r})}{16\pi t^4}(r\acute{r})[1-\cos(v+2k\beta\pi)]-\frac{(r-\acute{r})^3}{32\pi t^4}\\
&+\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{4\pi t^3}-\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}\\
&-\frac{1}{r}\frac{[(r\acute{r}[1-\cos(v+2k\beta\pi)])^2}{16\pi t^4}\}\\
+\;& e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{r}{4\pi t^3}E(\frac{r\acute{r}}{2t},v)-\frac{1}{2\pi\beta}\frac{r}{32\pi t^4}(r^2+\acute{r}^2)E(\frac{r\acute{r}}{2t},v)\\
&+\frac{1}{2\pi\beta}\frac{r(r\acute{r})}{16\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{r}\frac{r\acute{r}}{4\pi t^3}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\\
&+\frac{1}{2\pi\beta}\frac{1}{r}\frac{r\acute{r}}{8\pi t^2}\frac{r^2+\acute{r}^2}{4t^2}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{r}\frac{(r\acute{r})^2}{16\pi t^4}\frac{\partial^2E(\frac{r\acute{r}}{2t},v)}{\partial z^2}\}.
\end{aligned}$$

Then using (56), Theorem 10.1, (49), and (50), we get

$$\begin{aligned}
&\sup_{1\le t\le 2}|\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{(r-\acute{r})}{4\pi t^3}\\
&-\frac{(r-\acute{r})}{16\pi t^4}[r\acute{r}(1-\cos v)]-\frac{(r-\acute{r})^3}{32\pi t^4}\\
&+\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{4\pi t^3}-\frac{1}{r}\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^2}\frac{(r-\acute{r})^2}{4t^2}\\
&-\frac{1}{r}\frac{[(r\acute{r}[1-\cos(v+2k\beta\pi)])^2}{16\pi t^4}\}|\\
\le\;& Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}
\end{aligned}$$

and

$$\begin{aligned}
&\sup_{1\le t\le 2}|e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{r}{4\pi t^3}E(\frac{r\acute{r}}{2t},v)-\frac{1}{2\pi\beta}\frac{r}{32\pi t^4}(r^2+\acute{r}^2)E(\frac{r\acute{r}}{2t},v)\\
&+\frac{1}{2\pi\beta}\frac{r(r\acute{r})}{16\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{r}\frac{r\acute{r}}{4\pi t^3}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\\
&+\frac{1}{2\pi\beta}\frac{1}{r}\frac{r\acute{r}}{8\pi t^2}\frac{r^2+\acute{r}^2}{4t^2}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{1}{r}\frac{(r\acute{r})^2}{16\pi t^4}\frac{\partial^2 E(\frac{r\acute{r}}{2t},v)}{\partial z^2}\}|\\
\le\;& C(1+\acute{r})[1+(r^2+\acute{r}^2)^2]e^{-\frac{r^2+\acute{r}^2}{12}}\\
\le\;& Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Thus the estimate of $\sup_{1\le t\le 2}|\frac{\partial^2\widehat{H}}{\partial t\partial r}|$ follows. □

Compared to the previous ones, the next lemma seems to be a lot easier. Since there is no term concerning $\frac{1}{r}$.

Lemma 9.5. *We have*

$$\begin{aligned}
&\sup_{1\le t\le 2}|\frac{\partial\widehat{H}}{\partial t}|+\sup_{1\le t\le 2}|\frac{\partial^2\widehat{H}}{\partial t^2}|\\
\le\;& Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-\frac{r\acute{r}[1-\cos\beta(\theta-\theta')]}{8}}+Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}+Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Proof. of Lemma 9.4: We write $v=\beta(\theta-\theta')$. We still assume that $v\neq\pi$ or $-\pi$ mod $2\beta\pi$ so the last term in (19) vainishes. The conclusion for all v follows from continuity of the estimates with respect to v. The proof is more or less routine after applying the ideas in Lemma 9.3, but we want to emphasize the corresponding bound for each term here. We compute

$$
\begin{aligned}
&\frac{\partial \widehat{H}}{\partial t} \\
= \ & e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)}{4t}}\{-\frac{1}{4\pi t^2}+\frac{(r-\acute{r})^2}{16\pi t^3}+\frac{r\acute{r}(1-\cos v)}{8\pi t^3}\} \\
&+\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi} e^{-\frac{(r-\acute{r})^2}{4t}} e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}\{-\frac{1}{4\pi t^2} \\
&+\frac{(r-\acute{r})^2}{16\pi t^3}+\frac{r\acute{r}(1-\cos[v+2k\beta\pi])}{8\pi t^3}\} \\
&+e^{-\frac{r^2+\acute{r}^2}{4t}}\{-\frac{1}{2\pi\beta}\frac{1}{4\pi t^2}E(\frac{r\acute{r}}{2t},v)+\frac{1}{2\pi\beta}\frac{r^2+\acute{r}^2}{16\pi t^3}E(\frac{r\acute{r}}{2t},v) \\
&-\frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^3}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\}. \qquad (59)
\end{aligned}
$$

Then by the same idea as in the estimate of $\sup_{1\leq t\leq 2}|\frac{1}{r}\frac{\partial \widehat{H}}{\partial v}|$, using (56), (49), and (50), we get

$$
\begin{aligned}
&\sup_{1\leq t\leq 2}|\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi} e^{-\frac{(r-\acute{r})^2}{4t}} e^{-\frac{r\acute{r}}{2t}[1-\cos(v+2k\beta\pi)]}\{-\frac{1}{4\pi t^2} \\
&+\frac{(r-\acute{r})^2}{16\pi t^3}+\frac{r\acute{r}(1-\cos[v+2k\beta\pi])}{8\pi t^3}\}| \\
\leq \ & C(1+|r-\acute{r}|^2+r\acute{r})e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}} \\
\leq \ & Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}
\end{aligned}
$$

and

$$
\begin{aligned}
&\sup_{1\leq t\leq 2}|e^{-\frac{r^2+\acute{r}^2}{4t}}\{-\frac{1}{2\pi\beta}\frac{1}{4\pi t^2}E(\frac{r\acute{r}}{2t},v)+\frac{1}{2\pi\beta}\frac{r^2+\acute{r}^2}{16\pi t^3}E(\frac{r\acute{r}}{2t},v) \\
&-\frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^3}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\}| \\
\leq \ & C(1+r^2+\acute{r}^2)e^{-\frac{r^2+\acute{r}^2}{8}} \\
\leq \ & Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}
$$

thus the estimate for $\sup_{1\leq t\leq 2}|\frac{\partial \widehat{H}}{\partial t}|$ follows.

Continue differentiating (59) with respect to t we get

$$\begin{aligned}
&\frac{\partial^2 \widehat{H}}{\partial t^2}\\
= \ & e^{-\frac{(r-\acute{r})^2+2r\acute{r}(1-\cos v)}{4t}}\{\frac{1}{2\pi t^3} - \frac{(r-\acute{r})^2}{16\pi t^4} - \frac{r\acute{r}(1-\cos v)}{8\pi t^4} - \frac{3(r-\acute{r})^2}{16\pi t^4}\\
&+\frac{(r-\acute{r})^4}{64\pi t^5} + \frac{(r-\acute{r})^2 r\acute{r}(1-\cos v)}{32\pi t^5}\\
&-\frac{3r\acute{r}(1-\cos v)}{8\pi t^4} + \frac{[r\acute{r}(1-\cos v)]^2}{16\pi t^5} + \frac{(r-\acute{r})^2 r\acute{r}(1-\cos v)}{32\pi t^5}\}\\
&\\
+ \ & \Sigma_{k\neq 0,|v+2k\beta\pi|<\pi} e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{1}{2\pi t^3} - \frac{(r-\acute{r})^2}{16\pi t^4}\\
&-\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^4} - \frac{3(r-\acute{r})^2}{16\pi t^4}\\
+ \ & \frac{(r-\acute{r})^4}{64\pi t^5} + \frac{(r-\acute{r})^2 r\acute{r}[1-\cos(v+2k\beta\pi)]}{32\pi t^5} - \frac{3r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^4}\\
&+\frac{[r\acute{r}[1-\cos(v+2k\beta\pi)]]^2}{16\pi t^5} + \frac{(r-\acute{r})^2 r\acute{r}[1-\cos(v+2k\beta\pi)]}{32\pi t^5}\}\\
&\\
+ \ & e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{1}{2\pi t^3}E(\frac{r\acute{r}}{2t},v) - \frac{1}{2\pi\beta}\frac{r^2+\acute{r}^2}{16\pi t^4}E(\frac{r\acute{r}}{2t},v) + \frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\\
&-\frac{1}{2\pi\beta}\frac{3(r^2+\acute{r}^2)}{16\pi t^4}E(\frac{r\acute{r}}{2t},v) + \frac{1}{2\pi\beta}\frac{(r^2+\acute{r}^2)^2}{64\pi t^5}E(\frac{r\acute{r}}{2t},v)\\
&-\frac{1}{2\pi\beta}\frac{(r^2+\acute{r}^2)r\acute{r}}{32\pi t^5}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z} + \frac{1}{2\pi\beta}\frac{3r\acute{r}}{8\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\\
&-\frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^3}\frac{r^2+\acute{r}^2}{4t^2}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z} + \frac{1}{2\pi\beta}\frac{(r\acute{r})^2}{16\pi t^5}\frac{\partial^2 E(\frac{r\acute{r}}{2t},v)}{\partial z^2}\}.
\end{aligned}$$

Then using (56), Theorem 10.1, (49), (50), and the inequality

$$\{1+(r-\acute{r})^4+(r\acute{r})^2\}e^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}} \leq C,$$

we get

$$\begin{aligned}
&\sup_{1\le t\le 2}|\Sigma_{k\neq 0,|v+2k\beta\pi|<\pi}e^{-\frac{(r-\acute{r})^2+2r\acute{r}[1-\cos(v+2k\beta\pi)]}{4t}}\{\frac{1}{2\pi t^3}-\frac{(r-\acute{r})^2}{16\pi t^4}\\
&-\frac{r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^4}-\frac{3(r-\acute{r})^2}{16\pi t^4}\\
&+\frac{(r-\acute{r})^4}{64\pi t^5}+\frac{(r-\acute{r})^2r\acute{r}[1-\cos(v+2k\beta\pi)]}{32\pi t^5}-\frac{3r\acute{r}[1-\cos(v+2k\beta\pi)]}{8\pi t^4}\\
&+\frac{[r\acute{r}[1-\cos(v+2k\beta\pi)]]^2}{16\pi t^5}+\frac{(r-\acute{r})^2r\acute{r}[1-\cos(v+2k\beta\pi)]}{32\pi t^5}\}|\\
\le\ & C\{1+(r-\acute{r})^4+(r\acute{r})^2\}e^{-\frac{(r-\acute{r})^2}{8}}e^{-2ar\acute{r}}\\
\le\ & Ce^{-\frac{(r-\acute{r})^2}{16}}e^{-ar\acute{r}}
\end{aligned}$$

and

$$\begin{aligned}
&\sup_{1\le t\le 2}|e^{-\frac{r^2+\acute{r}^2}{4t}}\{\frac{1}{2\pi\beta}\frac{1}{2\pi t^3}E(\frac{r\acute{r}}{2t},v)-\frac{1}{2\pi\beta}\frac{r^2+\acute{r}^2}{16\pi t^4}E(\frac{r\acute{r}}{2t},v)\\
&+\frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}-\frac{1}{2\pi\beta}\frac{3(r^2+\acute{r}^2)}{16\pi t^4}E(\frac{r\acute{r}}{2t},v)+\frac{1}{2\pi\beta}\frac{(r^2+\acute{r}^2)^2}{64\pi t^5}E(\frac{r\acute{r}}{2t},v)\\
&-\frac{1}{2\pi\beta}\frac{(r^2+\acute{r}^2)r\acute{r}}{32\pi t^5}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}+\frac{1}{2\pi\beta}\frac{3r\acute{r}}{8\pi t^4}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}\\
&-\frac{1}{2\pi\beta}\frac{r\acute{r}}{8\pi t^3}\frac{r^2+\acute{r}^2}{4t}\frac{\partial E(\frac{r\acute{r}}{2t},v)}{\partial z}+\frac{1}{2\pi\beta}\frac{(r\acute{r})^2}{16\pi t^5}\frac{\partial^2 E(\frac{r\acute{r}}{2t},v)}{\partial z^2}\}|\\
\le\ & C[1+(r^2+\acute{r}^2)^2]e^{-\frac{r^2+\acute{r}^2}{8}}\\
\le\ & Ce^{-\frac{r^2+\acute{r}^2}{16}}.
\end{aligned}$$

Thus, the estimate of $\sup_{1\le t\le 2}|\frac{\partial^2\widehat{H}}{\partial t^2}|$ follows. □

10 Decay estimates for the heat kernel.

In this section, we prove the following theorem on the estimate of the E term in the heat kernel representation in Theorem 4.4.

Theorem 10.1. *For all $z\ge 0$ and v such that $v\neq\pi$ nor $-\pi$ mod $2\beta\pi$, the function $E(z,v)$ in Theorem 4.4 satisfies the following estimates.*

- *For any $p\ge n-1,\ n\ge 0$, there exists a $C_{p,n}$ such that*

$$|\frac{\partial}{\partial v}\{z^p\frac{\partial^n}{\partial z^n}E(z,v)\}|\le C_{p,n};$$

- *For any* $p \geq \max\{n-1, 0\}$, $n \geq 0$, *there exists a* $\widehat{C}_{p,n}$ *such that*
$$|z^p \frac{\partial^n}{\partial z^n} E(z,v)\}| \leq \widehat{C}_{p,n}.$$

Remark 10.2. The points $v = \pi$ or $-\pi$ *mod* $2\beta\pi$ are the discontiuities of $E(z,v)$. Our theorem shows that $E(z,v)$ behaves very well away from these discontinuity points of v, which is sufficient to apply.

Proof. of Theorem 10.1: By using inductions and inequality 50, these estimates are directly corollaries of Lemma 10.3 and Lemma 10.4 below. ☐

Lemma 10.3. *For any nonnegative integer* $n \geq 1$, *there exists a* C_n *with the following property. For all* $z \geq 0$, v *such that* $v \neq \pi$ *nor* $-\pi$ *mod* $2\beta\pi$, *the following estimates hold:*
$$|z^{n-1} \frac{\partial^n}{\partial z^n}[e^z E(z,v)]| \leq C_n,$$
and
$$|e^z E(z,v)| \leq C.$$

Proof. of Lemma 10.3: Actually the estimates also hold when $v = \pi$ or $-\pi$ *mod* $2\beta\pi$. However, since $E(z,v)$ is not continuous when $v = \pi$ or $-\pi$ *mod* $2\beta\pi$, we want to concentrate on the estimates close to these discontinuity points, but not exactly at these points. When $n \geq 1$, we compute from Theorem 4.4 that
$$\begin{aligned} & z^{n-1}\frac{d^n[e^z E(z,v)]}{dz^n} \\ = & \int_0^\infty (-z(\cosh y - 1))^{n-1} e^{-z(\cosh y - 1)}(1-\cosh y) \\ & \frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta} - \cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]} dy. \end{aligned} \tag{60}$$

To prove $|z^{n-1}\frac{d^n[e^z E(z,v)]}{dz^n}| \leq C_n$, we divide the situation into two cases below.

Case 1, suppose $\frac{1}{\beta}$ is an integer, then $\sin\frac{\pi}{\beta} = 0$. Then $E(z,v) \equiv 0$, the conclusion of Lemma 10.3 immediately follows.

Case 2, suppose $\frac{1}{\beta}$ is not an integer. Then there is a C such that $|\cos\frac{v-\pi}{\beta} - 1| \leq \frac{1}{C}$ implies $|\cos\frac{v+\pi}{\beta} - 1| \geq \frac{1}{C}$, and $|\cos\frac{v+\pi}{\beta} - 1| \leq \frac{1}{C}$ implies $|\cos\frac{v-\pi}{\beta} - 1| \geq \frac{1}{C}$.

Then the only issue we should worry about is that when $\cos\frac{v+\pi}{\beta}$ (or $\cos\frac{v-\pi}{\beta}$) is very close to 1, and y is close to 0, the integral (60) might not be uniformly bounded with respect to v. We will show that this bad situation won't happen. Notice that if both $|\cos\frac{v-\pi}{\beta}-1|\geq\frac{1}{C}$ and $|\cos\frac{v+\pi}{\beta}-1|\geq\frac{1}{C}$ hold, we trivially have $|z^{n-1}\frac{d^n[e^zE(z,v)]}{dz^n}|\leq C_n$.

Thus it suffices to assume $|\cos\frac{v-\pi}{\beta}-1|\leq\frac{1}{C}$ and consequently $|\cos\frac{v+\pi}{\beta}-1|\geq\frac{1}{C}$. The case when $|\cos\frac{v+\pi}{\beta}-1|\leq\frac{1}{C}$ is the same.

First we compute

$$\frac{\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{y}{\beta}}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}=\frac{\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}-\cos\frac{v}{\beta}, \tag{61}$$

thus our primary interest is in the term $\frac{\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}}{\cosh\frac{y}{\beta}}$. Then using the inequality

$$\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}\geq\frac{y^2}{2\beta^2}+2\sin^2\frac{v-\pi}{2\beta},$$

we have for any positive real number A that

$$\begin{aligned}
&|\int_0^A\frac{\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}dy|\\
\leq\ & 2\beta^2|\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}|\int_0^A\frac{1}{y^2+\beta^2\sin^2\frac{v-\pi}{2\beta}}dy\\
=\ & 2\beta^2|\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}|\times\frac{\arctan\frac{A}{\beta|\sin\frac{v-\pi}{2\beta}|}}{\beta|\sin\frac{v-\pi}{2\beta}|}\\
=\ & \frac{2\beta|\cos\frac{\pi}{\beta}\sin^2\frac{v-\pi}{\beta}+\sin\frac{\pi}{\beta}\cos\frac{v-\pi}{\beta}\sin\frac{v-\pi}{\beta}|}{|\sin\frac{v-\pi}{2\beta}|}\times\arctan\frac{A}{\beta|\sin\frac{v-\pi}{2\beta}|}\\
\leq\ & 4\pi\beta.
\end{aligned}\tag{62}$$

By the same method, we have for any A that

$$\int_0^A\frac{1}{\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}}dy\leq\beta^2\int_0^A\frac{1}{y^2+\frac{\beta^2}{C}}dy=\sqrt{C}\beta\arctan\frac{A\sqrt{C}}{\beta}\leq\sqrt{C}\beta\frac{\pi}{2}.$$

It's easy to deduce the following inequality.

$$|z^{n-1}\frac{d^ne^{-z(\cosh y-1)}}{dz^n}|\leq C_n(\cosh y-1). \tag{63}$$

Furthurmore, Using the inequality $|\cos\frac{v+\pi}{\beta}-1|\geq\frac{1}{C}$ and equation (61), it's easy to see that

$$\begin{aligned}
&|\frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]}| \\
\leq\ & C|\frac{\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{v-\pi}{\beta}}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}+\frac{1}{\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}}|. \qquad (64)
\end{aligned}$$

Then using (63) and (64), we deduce

$$\begin{aligned}
&|z^{n-1}\frac{d^n e^z E(z,v)}{dz^n}| \\
=\ & |\int_0^\infty z^{n-1}\frac{d^n e^{-z(\cosh y-1)}}{dz^n}\frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]}dy| \quad (65) \\
\leq\ & C_n\int_0^\infty \frac{(\cosh y-1)|\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}|}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}+\frac{(\cosh y-1)}{\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}}dy.
\end{aligned}$$

Since $\frac{1}{\beta}>1$ and $|\cos\frac{v+\pi}{\beta}-1|\geq\frac{1}{C}$, we have

$$\int_0^\infty\frac{\cosh y-1}{\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}}dy\leq C. \qquad (66)$$

Using almost the same trick as in (62) we make the following estimate:

$$\begin{aligned}
&\int_0^\infty\frac{(\cosh y-1)|\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}|}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}dy \\
\leq\ & 5\int_0^1\frac{|\cos\frac{\pi}{\beta}-\cos\frac{v}{\beta}\cos\frac{v-\pi}{\beta}|}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}dy+3\int_1^\infty\frac{\cosh y-1}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}dy \\
\leq\ & C. \qquad (67)
\end{aligned}$$

Thus, combining (67), (66), and (65), we conclude that

$$|z^{n-1}\frac{\partial^n}{\partial z^n}[e^z E(z,v)]|\leq C_n.$$

The conclusion $|e^z E(z,v)|\leq C$ is obvious from the discussion above. ☐

Lemma 10.4. *For any nonnegative integer $n\geq 0$, there exists a C_n with the following property. For all $z\geq 0$, v such that $v\neq\pi$ nor $-\pi$ mod $2\beta\pi$, the following estimate holds*

$$|\frac{\partial}{\partial v}\{z^{n-1}\frac{\partial^n}{\partial z^n}[e^z E(z,v)]\}|\leq C_n.$$

Proof. of Lemma 10.4: The proof here is more complicated than that of Lemma 10.3. We only prove the statement for $|\frac{\partial^2}{\partial v \partial z} e^z E(z,v)|$, the proofs of $|\frac{\partial}{\partial v}[z^{n-1}\frac{\partial}{\partial z^n} e^z E(z,v)]|$ are by the same way and even easier. Recall that

$$E(z,v) = \int_0^\infty e^{-z\cosh y} F(v,y) dy,$$

where

$$F(v,y) = \frac{2\sin\frac{\pi}{\beta}[\cos\frac{\pi}{\beta} - \cos\frac{v}{\beta}\cosh\frac{y}{\beta}]}{[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}][\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]}.$$

We compute

$$\begin{aligned} & \frac{\partial}{\partial v} F(v,y) \\ = \quad & \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}[\cosh^3\frac{y}{\beta} - (2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\cosh\frac{y}{\beta} + \cos\frac{v+\pi}{\beta} + \cos\frac{v-\pi}{\beta}]}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}]^2[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2}. \end{aligned}$$

Therefore we have the following integral representation of $\frac{\partial^2}{\partial v \partial z}[e^z E(z,v)]$:

$$\begin{aligned} & \frac{\partial^2}{\partial v \partial z} e^z E(z,v) \\ = \quad & \int_0^\infty \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}[\cosh^3\frac{y}{\beta} - (2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\cosh\frac{y}{\beta} + \cos\frac{v+\pi}{\beta} + \cos\frac{v-\pi}{\beta}]}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}]^2[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2} \\ & -(\cosh y - 1)e^{-z(\cosh y - 1)} dy. \end{aligned} \tag{68}$$

With the same idea as in Lemma 10.3 in mind, we divide the situation into two cases below.

Case 1, suppose $\frac{1}{\beta}$ is an integer, then $\sin\frac{\pi}{\beta} = 0$. Then $E(z,v) \equiv 0$.

Case 2, suppose $\frac{1}{\beta}$ is not an integer. Then there is a C such that $|\cos\frac{v-\pi}{\beta} - 1| \le \frac{1}{C}$ implies $|\cos\frac{v+\pi}{\beta} - 1| \ge \frac{1}{C}$, and $|\cos\frac{v+\pi}{\beta} - 1| \le \frac{1}{C}$ implies $|\cos\frac{v-\pi}{\beta} - 1| \ge \frac{1}{C}$.

Again, the only issue we should worry about is that when $\cos\frac{v+\pi}{\beta}$ or $\cos\frac{v-\pi}{\beta}$ is very close to 1, and y is close to 0, the integral (71) might not be uniformly bounded with respect to v. We will show this bad situation won't happen. Notice that if both $|\cos\frac{v-\pi}{\beta} - 1| \ge \frac{1}{C}$ and $|\cos\frac{v+\pi}{\beta} - 1| \ge \frac{1}{C}$ hold, we trivially have $|\frac{\partial^2}{\partial v \partial z} e^z E(z,v)| \le C$.

Thus it suffices to assume $|\cos\frac{v-\pi}{\beta}-1|\le\frac{1}{C}$ and consequently $|\cos\frac{v+\pi}{\beta}-1|\ge\frac{1}{C}$. The case when $|\cos\frac{v+\pi}{\beta}-1|\le\frac{1}{C}$ is the same.

Next we start discussing on the case $|\cos\frac{v-\pi}{\beta}-1|\le\frac{1}{C}$. It is easy to see that

$$\begin{aligned}&\frac{\cosh^3\frac{y}{\beta}-(2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\cosh\frac{y}{\beta}+\cos\frac{v+\pi}{\beta}+\cos\frac{v-\pi}{\beta}}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}]^2}\\ = \; &\cosh\frac{y}{\beta}+2\cos\frac{v-\pi}{\beta}+\frac{3\cos^2\frac{v-\pi}{\beta}-\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta}-2}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}\\ &-\frac{\sin^2\frac{v-\pi}{\beta}[\cos\frac{v-\pi}{\beta}-\cos\frac{v+\pi}{\beta}]}{(\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta})^2}.\end{aligned}$$

Denote

$$S(z,y)=\frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}(1-\cosh y)e^{-z(\cosh y-1)}}{\beta[\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]^2},$$

$$D_1(v,y)=\frac{1}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}-\frac{1}{\frac{y^2}{2\beta^2}+1-\cos\frac{v-\pi}{\beta}}.$$

Then we have the following splitting

$$\begin{aligned}&\frac{\partial^2}{\partial v\partial z}e^zE(z,v)\\ = \; &\int_0^{+\infty}S(z,y)\frac{\cosh^3\frac{y}{\beta}-(2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\frac{y}{\beta}+\cos\frac{v+\pi}{\beta}+\cos\frac{v-\pi}{\beta}}{[\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}]^2}dy\\ = \; &\int_1^{+\infty}(1-\cosh y)e^{-z(\cosh y-1)}\frac{\partial F(y,v)}{\partial v}dy+\int_0^1S(z,y)(\cosh\frac{y}{\beta}\\ &+2\cos\frac{v-\pi}{\beta})dy+\int_0^1S(z,y)\frac{3\cos^2\frac{v-\pi}{\beta}-\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta}-2}{\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta}}dy\\ &-\int_0^1S(z,y)\frac{\sin^2\frac{v-\pi}{\beta}[\cos\frac{v-\pi}{\beta}-\cos\frac{v+\pi}{\beta}]}{(\cosh\frac{y}{\beta}-\cos\frac{v-\pi}{\beta})^2}dy\\ = \; &\Sigma_{k=1}^7II_k.\end{aligned}$$

The terms in the splitting are

$$
\begin{aligned}
II_1 &= \int_1^{+\infty} -(\cosh y - 1)e^{-z(\cosh y-1)}\frac{\partial F(y,v)}{\partial v}dy,\\
II_2 &= \int_0^1 S(z,y)(\cosh\frac{y}{\beta} + 2\cos\frac{v-\pi}{\beta})dy,\\
II_3 &= (3\cos^2\frac{v-\pi}{\beta} - \cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta} - 2)\int_0^1 S(z,y)D_1(v,y)dy,\\
II_4 &= (3\cos^2\frac{v-\pi}{\beta} - \cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta} - 2)\int_0^1 \frac{S(z,y)}{\frac{y^2}{2\beta^2} + 1 - \cos\frac{v-\pi}{\beta}}dy,\\
II_5 &= -\sin^2\frac{v-\pi}{\beta}[\cos\frac{v-\pi}{\beta} - \cos\frac{v+\pi}{\beta}]\int_0^1 S(z,y)D_1^2(v,y)dy,\\
II_6 &= -2\sin^2\frac{v-\pi}{\beta}[\cos\frac{v-\pi}{\beta} - \cos\frac{v+\pi}{\beta}]\int_0^1 S(z,y)\frac{D_1(v,y)}{\frac{y^2}{2\beta^2} + 1 - \cos\frac{v-\pi}{\beta}}dy,\\
II_7 &= -\sin^2\frac{v-\pi}{\beta}[\cos\frac{v-\pi}{\beta} - \cos\frac{v+\pi}{\beta}]\int_0^1 \frac{S(z,y)}{|\frac{y^2}{2\beta^2} + 1 - \cos\frac{v-\pi}{\beta}|^2}dy.
\end{aligned}
$$

We have the following claim regarding D_1 and S.

Claim 10.5. *There exists a C such that when $y \leq 1$, we have*

$$|D_1(v,y)| \leq C, \quad |S(z,y)| \leq Cy^2.$$

The assumption $y \leq 1$ is important. It's easy to see $|S(z,y)| \leq Cy^2$ from the inequality $|\cos\frac{v+\pi}{\beta} - 1| \geq \frac{1}{C}$.

The estimate $|D_1(v,y)| \leq C$ follows from the following inequality

$$D_1(v,y) = \frac{O(y^4)}{(\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta})(\frac{y^2}{2\beta^2} + 1 - \cos\frac{v-\pi}{\beta})} \leq \frac{O(y^4)}{y^4} \leq C$$

when $y \leq 1$, thanks to the fact that the Taylor series of $\cosh y$ at 0 only contain even order terms. Then Claim 10.5 is proved.

From Claim 10.5 we immediately see

$$|II_2| + |II_3| + |II_4| + |II_5| + |II_6| \leq C, \tag{69}$$

and

$$
\begin{aligned}
|II_7| &\leq C\int_0^1 \frac{y^2\sin^2(\frac{v-\pi}{\beta})}{|\frac{y^2}{2\beta^2} + 2\sin^2\frac{v-\pi}{2\beta}|^2}dy\\
&\leq C. \qquad (70)
\end{aligned}
$$

Next we show that II_1 could also be easily bounded.

Claim 10.6. $|II_1| < C$.

Claim 10.6 follows from the trivial fact that $\frac{1}{\beta} > 1$ and the following estimates when $y \geq 1$.

$$\begin{aligned} |(-(\cosh y - 1))e^{-z(\cosh y-1)}\frac{\partial}{\partial v}F(v,y)| &\leq C\frac{(\cosh y - 1)(\cosh^3 \frac{y}{\beta})}{\cosh^4 \frac{y}{\beta}} \\ &\leq Ce^{-(\frac{1}{\beta}-1)y}. \end{aligned}$$

Since the last term $Ce^{-(\frac{1}{\beta}-1)y}$ is integrable, then Claim 10.6 follows.

Then when $n = 1$, Lemma 10.4 follows from (69), (70), claim 10.6.

When $n = 0$, there is a point we should clearify. With respect to $\frac{1}{z}\frac{\partial}{\partial v}e^z E(z,v)$, we apply the following fact which is straight forward from Theorem 4:

$$\frac{\partial}{\partial v}E(0,v) = 0 \text{ when } v \neq \ \pi \text{ nor} - \pi \text{ mod } 2\beta\pi. \tag{71}$$

Notice that $E(z,v)$ is not differentiable when $v = \pi$ or $-\pi$ *mod* $2\beta\pi$.

The fact in (71) means that when $v = \pi$ or $-\pi$ *mod* $2\beta\pi$, we have

$$\begin{aligned} &\textstyle\int_0^\infty \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}[\cosh^3\frac{y}{\beta} - (2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\cosh\frac{y}{\beta} + \cos\frac{v+\pi}{\beta} + \cos\frac{v-\pi}{\beta}]}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}]^2[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2}dy \\ = &\ 0. \end{aligned}$$

Then

$$\begin{aligned} &\frac{1}{z}\frac{\partial}{\partial v}e^z E(z,v) \\ = &\int_0^\infty \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}[\cosh^3\frac{y}{\beta} - (2+\cos\frac{v-\pi}{\beta}\cos\frac{v+\pi}{\beta})\cosh\frac{y}{\beta} + \cos\frac{v+\pi}{\beta} + \cos\frac{v-\pi}{\beta}]}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v-\pi}{\beta}]^2[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2} \\ \times &\ \frac{e^{-z(\cosh y-1)} - 1}{z}dy. \end{aligned}$$

Then instead of considering $S(z,y) = \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}(1-\cosh y)e^{-z(\cosh y-1)}}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2}$ (in the case of $\frac{\partial^2}{\partial v\partial z}e^z E(z,v)$), we consider

$$\widehat{S}(z,y) = \frac{2\sin\frac{\pi}{\beta}\sin\frac{v}{\beta}(e^{-z(\cosh y-1)} - 1)}{\beta[\cosh\frac{y}{\beta} - \cos\frac{v+\pi}{\beta}]^2 z}$$

in the case of $\frac{\partial}{\partial v}e^zE(z,v)$. Apparently when $v\neq\pi$ nor $-\pi$ mod $2\beta\pi$, we still have $\frac{1}{[\cosh\frac{y}{\beta}-\cos\frac{v+\pi}{\beta}]^2}\leq C$; when and $y\leq 1$, we still have that

$$\widehat{S}(z,y)\leq Cy^2.$$

Therefore the proof for $\frac{\partial^2}{\partial v\partial z}e^zE(z,v)$ exactly carries over to the proof of $\frac{\partial}{\partial v}e^zE(z,v)$ line by line. $\square$

11 Appendix: some lower order estimates.

In this appendix, we enumerate some interpolation formulas needed for the Schauder estimate. We also state some lower order Schauder estimates. As in [20], they are necessary but routine and simple. Some of them are not proved by the same way as in [20]. We would like to point out that comparing to the proof of proposition 5.1(which is the top order seminorm estimate and is our main work in this article,) the following lower order estimates are really similar and much much easier.

Lemma 11.1. *For any $\epsilon>0$ and domain Ω, the following parabolic interpolation inequalities hold.*

$$\begin{cases} [\nabla u]^{(1)}_{0,\Omega\times[0,T]} & \leq\ C(\epsilon)|u|_{0,\Omega\times[0,T]}+\epsilon[\nabla u]^{(1)}_{\alpha,\beta,\Omega\times[0,T]};\\ [u]^{(*)}_{\alpha,\beta,\Omega\times[0,T]} & \leq\ \epsilon[\nabla u]^{(1)}_{0,\Omega\times[0,T]}+C(\epsilon)[u]_{0,\Omega\times[0,T]};\\ |\Delta u|^{(2)}_{0,\Omega\times[0,T]} & \leq\ C(\epsilon)|\nabla u|^{(1)}_{0,\Omega\times[0,T]}+\epsilon|\Delta u|^{(2)}_{\alpha,\beta,\Omega\times[0,T]};\\ |\sqrt{-1}\partial\bar{\partial}u|^{(2)}_{0,\Omega\times[0,T]} & \leq\ C(\epsilon)|\nabla u|^{(1)}_{0,\Omega\times[0,T]}+\epsilon|\sqrt{-1}\partial\bar{\partial}u|^{(2)}_{\alpha,\beta,\Omega\times[0,T]}. \end{cases}$$

Proof. of Lemma 11.1: The second item is by standard calculations in [20]. We need to point out that, on the first inequality

$$[\nabla u]^{(1)}_{0,\Omega\times[0,T]}\leq C(\epsilon)|u|_{0,\Omega\times[0,T]}+\epsilon[\nabla u]^{(1)}_{\alpha,\beta,\Omega\times[0,T]},$$

we have to deal with the term $\frac{1}{r}\frac{\partial u}{\partial\theta}$, in a different way. And the observation is that, for any $\widehat{x}$ and $r_0>0$, over every level surface $r=r_0$, there is a θ_0, such that $\frac{\partial u}{\partial\theta}(r_0,\theta_0,\widehat{x})=0$.

The third item concerns integration by parts. The fourth item concern integration by parts over holomorphic disks. The rest of the proof are regular as in . $\square$

Similarly, using parabolic weights we can prove the following intepolation inequalities, which are applied also in the proof of Theorem 1.8 in section 5.

Lemma 11.2. *For any $\epsilon > 0$ and domain Ω, the following parabolic interpolation inequalities hold.*

$$\begin{cases} [\nabla u]^{[1]}_{0,\Omega\times[0,T]} & \leq \; C(\epsilon)|u|_{0,\Omega\times[0,T]} + \epsilon[\nabla u]^{(1)}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]}; \\ [u]^{[*]}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]} & \leq \; \epsilon[\nabla u]^{[1]}_{0,\Omega\times[0,T]} + C(\epsilon)[u]_{0,\Omega\times[0,T]}; \\ |\Delta u|^{[2]}_{0,\Omega\times[0,T]} & \leq \; C(\epsilon)|\nabla u|^{[1]}_{0,\Omega\times[0,T]} + \epsilon|\Delta u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]}; \\ |\sqrt{-1}\partial\bar{\partial}u|^{[2]}_{0,\Omega\times[0,T]} & \leq \; C(\epsilon)|\nabla u|^{[1]}_{0,\Omega\times[0,T]} + \epsilon|\sqrt{-1}\partial\bar{\partial}u|^{[2]}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]}. \end{cases}$$

Apply the techniques in the previous sections we have the following lower order Schauder estimates.

Lemma 11.3. *Suppose $\alpha < (\min\frac{1}{\beta} - 1, 1)$. The following lower order Schauder estimates for the heat operator are true for all $u \in C^{2+\alpha,1+\frac{\alpha}{2},\beta}(\Omega\times[0,T])$ with zero initial value.*

$$\begin{cases} |i\partial\bar{\partial}u|^{(2)}_{0,\Omega\times[0,T]} & \leq \; C(\alpha)|u|_{0,\Omega\times[0,T]} + C(\alpha)|(\Delta - \frac{\partial}{\partial t})u|^{(2)}_{\alpha,\beta,\Omega\times[0,T]}; \\ |\frac{\partial u}{\partial t}|^{(2)}_{0,\Omega\times[0,T]} & \leq \; C(\alpha)|u|_{0,\Omega\times[0,T]} + C(\alpha)|(\Delta - \frac{\partial}{\partial t})u|^{(2)}_{\alpha,\beta,\Omega\times[0,T]}; \\ |\nabla u|^{(1)}_{\alpha,\frac{\alpha}{2},\beta,\Omega\times[0,T]} & \leq \; C_4|u|_{0,\Omega\times[0,T]} + C_4|(\Delta - \frac{\partial}{\partial t})u|^{(2)}_{0,\Omega\times[0,T]}. \end{cases}$$

The next theorem is about the necessary C^0 estimate. The proof of it only involves the parabolic maximal principle, which works in our conical $C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$ setting. We refer the readers to [15] for a full proof. The idea originates from Jeffres' work on the elliptic case in [22].

Lemma 11.4. *Suppose $\{a(t),\ t \in [0,T]\}$ is a $C^{\alpha,\frac{\alpha}{2},\beta}[0,T]$ family of (α,β) metrics (See Definition 2.5). Suppose $u \in C^{2+\alpha,1+\frac{\alpha}{2},\beta}[0,T]$ solves the following parabolic equation.*

$$\frac{\partial u}{\partial t} = \Delta_{a(t)}u + v,\ \ u(0,x) = 0. \tag{72}$$

Then the following estimate holds.

$$|u|_{0,M\times[0,T]} \leq C * T * |v|_{0,M\times[0,T]}.$$

References

[1] R, Berman. *A thermodynamic formalism for Monge-Ampere equations.* arXiv1011.3976.

[2] R, Berman. *K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics.* arxiv1205.6214.

[3] S, Brendle. *Ricci flat Kähler metrics with edge singularities.* To appear in International Mathematics Research Notices.

[4] S, Calamai; K, Zheng. *Geodesics in the space of Kähler cone metrics.* arXiv1205.0056.

[5] F, Campana; H, Guenancia; M, Paun. *Metrics with cone singularities along normal crossing divisors and holomorphic tensor fields .*

[6] H-D, Cao. *Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds.* Invent.Math.81(1985), no.2, 359-372.

[7] H,S, Carslaw. *The Green's Function For a Wedge of Any Angle, and Other Problems in the Conduction of Heat.*

[8] X-X, Chen; W-Y, Ding. *Ricci flow on surfaces with degenerate initial metrics.* J. Partial Differential Equations 20 (2007), no. 3.

[9] X-X, Chen; S, Donaldson; S, Sun. *Kähler-Einstein metrics and stability.* arXiv: 1210.7494. To appear in Int. Math. Res. Not (2013).

[10] X-X, Chen; S, Donaldson; S, Sun. *Kähler-Einstein metric on Fano manifolds, I: approximation of metrics with cone singularities.* arXiv1211.4566.

[11] X-X, Chen; S, Donaldson; S, Sun. *Kähler-Einstein metric on Fano manifolds, II: limits with cone angle less than* 2π. arXiv1212.4714.

[12] X-X, Chen; S, Donaldson; S, Sun. *Kähler-Einstein metric on Fano manifolds, III: limits with cone angle approaches* 2π *and completion of the main proof.* arXiv1302.0282.

[13] X-X, Chen; B, Wang. *Kähler-Ricci flow on Fano manifolds (I).* preprint. arXiv0909.2391.

[14] X-X, Chen; G, Tian; Z, Zhang. *On the weak Kähler-Ricci flow.* Trans. Amer. Math. Soc. 363 (2011), no. 6, 2849–2863.

[15] X-X, Chen; Y, Wang. *On the long time behavior of the Conical Kähler-Ricci flow.* Unpublished work.

[16] Donaldson, S: Scalar curvature and stability of toric varieties. J. Diff. Geom., **62** (2002), 289-349.

[17] S, Donaldson. *Kähler metrics with cone singularities along a divisor.*

[18] S, Donaldson. *Discussion of the Kahler-Einstein problem.* www.imperial.ac.uk/ skdona/KENOTES.PDF.

[19] P, Eyssidieux; V, Guedj; A, Zeriahi. *Singular Kähler-Einstein metrics.* J. Amer. Math. Soc. 22 (2009), 607-639.

[20] D,Gilbarg; N ,S, Trudinger. *Elliptic Partial Differential Equations of Second Order.* Springer.

[21] A. Hugues, *The space of Poincaré type Kähler metrics on the complement of a divisor.* eprint arXiv:1109.3159.

[22] T, Jeffres. *Uniqueness of Kähler-Einstein cone metrics*, Publ. Math. 44 (2000).

[23] J, Isenberg; R, Mazzeo; N, Sesum. *Ricci flow in two dimensions.* arxiv1103.4669.

[24] Jeffres, T; Mazzeo; R, Rubinstein. *Kähler-Einstein metrics with edge singularities.* To appear in Annals of Math.

[25] Ladyzenskaja; Solonnikov; Ural'ceva. *Linear and quasi-linear equations of parabolic type.* Translations of Mathematical Monographs 23, Providence, RI: American Mathematical Society.

[26] C, Li; S. Sun. *Conical Kähler-Einstein metric revisited.* arXiv1207.5011.

[27] G, Lieberman. *Second order parabolic equations.* World Scientific, 1996.

[28] F, Luo; G, Tian. *Liouville Equation and Spherical Convex Polytopes.* Proceedings of the American Mathematical Society, Vol. 116, No. 4 (Dec., 1992), pp.1119-1129.

[29] R, McOwen. *Point singularities and conformal metrics on Riemann surfaces.* Proc. Amer. Math. Soc. 103 (1988), 222-224.

[30] J, Möllers. *Heat kernel analysis for Bessel operators on symmetric cones.* arXiv:1209.2310.

[31] E, Mooers. *The heat kernel for manifolds with conic singularities.* PHD Thesis, MIT.

[32] D, Ramos. *Smoothening cone points with Ricci flow.* arxiv1109.5554.

[33] G, Székelyhidi; V, Tosatti. *Regularity of weak solutions of a complex Monge-Ampere equation.* Anal. PDE 4 (2011), no. 3.

[34] J, Song; X, Wang. *The greatest Ricci lower bound, conical Einstein metrics and the Chern number inequality.* arXiv1207.4839.

[35] J, Stoppa. *K-stability of constant scalar curvature Kähler manifolds.* Adv. Math. 221 (2009), no. 4, 13971408.

[36] S, Sun; Y, Wang . *On the Kähler-Ricci flow near a Kähler-Einstein metric.* To appear in J. Reine Angew. Math.

[37] T, Richard. *Canonical smoothing of compact Alexandrov surfaces via Ricci flow.* arXiv1204.5461.

[38] G, Tian. *On Kahler-Einstein metrics on complex surfaces with* $C_1 > 0$. Invent. Math. 89 (1987), no. 2, 225-246.

[39] G, Tian. *On Calabis conjecture for complex surfaces with positive first Chern class.* Invent. Math. 101 (1990), no.1, 101-172.

[40] G, Tian. *Kahler-Einstein metrics with positive scalar curvature.* Inv. Math. 130(1997), 1-39.

[41] G, Tian; S-T, Yau. *Complete Kähler Manifolds with Zero Ricci Curvature. I.* Journal of the American Mathematical Society, Vol. 3, No. 3. (Jul., 1990), pp. 579-609.

[42] G, Tian; S-T, Yau. *Kähler-Einstein metrics on complex surfaces with* $C_1 > 0$. Comm. Math. Phys. 112(1987), no. 1, 175-203.

[43] G, Tian; X-H, Zhu. *Convergence of Kähler-Ricci flow.* J. Amer. Math. Soc. 20 (2007), no. 3, 675–699.

[44] G, Tian; X-H, Zhu. *Perelman's W-functional and stability of Kähler-Ricci flow*, arxiv0801.3504.

[45] G, Tian; X-H, Zhu. *Stability of Kähler-Ricci flow on a Fano manifold II.* preprint.

[46] M, Troyanov. *Prescribing curvature on compact surfaces with conic singularities.* Trans. Amer. Math. Soc. 324 (1991).

[47] P, Topping. *Ricci flow compactness via pseudolocality, and flows with incomplete initial metrics.* J.E.M.S., 12 (2010) 1429–1451.

[48] Y, Wang. *Notes on the* L^2*-estimates and regularity of parabolic equations over conical manifolds.* Unpublished work.

[49] G,N, Watson. *A Treatise on the Theory of Bessel Functions*,Cambridge Mathematical Library.

[50] G,N, Watson; E,T, Whittaker. *A Course of Modern Analysis.* Cambridge University Press.

[51] S-T, Yau. *On the Ricci curvature of a compact Kähler manifold and the Complex Monge Amp'ere equation I.* Comm. Pure Appl. Math. 31 (1978).

[52] H, Yin. *Ricci flow on surfaces with conical singularities.* Journal of Geometric Analysis October 2010, Volume 20, Issue 4, pp 970-995.

Xiuxiong chen, Department of Mathematics, Stony Brook University, NY, USA; xiu@math.sunysb.edu.

Yuanqi Wang, Department of Mathematics, University of California at Santa Barbara, Santa Barbara, CA, USA; wangyuanqi@math.ucsb.edu.